

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number  
**WO 01/18225 A1**

(51) International Patent Classification<sup>7</sup>: C12N 15/90, A01K 67/027

(21) International Application Number: PCT/US99/30078

(22) International Filing Date:  
16 December 1999 (16.12.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/152,522 3 September 1999 (03.09.1999) US

(71) Applicant: XENOGEN CORPORATION [US/US]; 860 Atlantic Avenue, Alameda, CA 94501 (US).

(72) Inventor: ZHANG, Ning; Xenogen Corporation, 860 Atlantic Avenue, Alameda, CA 94501 (US).

(74) Agent: SHOLTZ, Charles, K.; Xenogen Corporation, 860 Atlantic Avenue, Alameda, CA 94501 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED THEREWITH



(57) Abstract: The present invention teaches targeting constructs and methods of use thereof for creating transgenic animals in which at least one single-copy, non-essential gene is replaced with a reporter expression cassette, for example, a luciferase gene operably linked to a promoter heterologous to the single-copy, non-essential gene. Thus, the present invention provides novel methods and vector constructs useful for the generation of transgenic animals. The invention further includes methods of using these animals.

## TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED THEREWITH

5

### TECHNICAL FIELD

This invention is in the field of molecular biology and medicine. More specifically, it relates to novel vector constructs and methods of use thereof for introducing heterologous polynucleotides into a host cell. Further, the invention relates to vector constructs and methods of use thereof to generate transgenic organisms, particularly transgenic mice.

### BACKGROUND

In recent years, mouse geneticists have succeeded in creating transgenic animals by manipulating the genes of developing embryos and introducing foreign genes into these embryos. Once these genes have integrated into the genome of the recipient embryo, the resulting embryos or adult animals can be analyzed to determine the function of the gene.

Traditionally, transgenic mice have been generated through DNA microinjection approach. Such an approach leads to the creation of "founder" mice having at least one copy of the transgene randomly integrated into the genome. Because neither the copy number nor the integration sites can be controlled, transgenic mice generated by this method are genetically different from each other. The expression of transgenes in such transgenic mice are not uniform because of, for example, the difference in copy numbers of the transgene. Furthermore, chromosomal location of the transgene often affects the expression level of the transgene. For most of *in vivo* studies, particularly ones where it is desirable to compare levels of gene expression across different animals, it is important to have the mice with little or no genetic variation (i.e., isogenic mice) in order to reduce the systematic error.

One way of controlling copy number and the chromosomal integration site is by using targeting constructs to create transgenic animals. U.S. Patent Nos. 5,464,764 and 5,487,992 describe this type of transgenic animal in which a gene of interest is deleted or mutated sufficiently to disrupt its function. These "knock-out" animals are made by taking advantage of the phenomena of homologous recombination. (See, also U.S.

Patent Nos. 5,631,153 and 5,627,059). Using the "knock-out" approach, when the vector is introduced into the embryonic stem cell, a sequence becomes integrated into a target gene in the genome via homologous recombination. The integration disrupts the function of the target gene allows for examination of the phenotype resulting from the  
5 disruption of the gene. Neither of the above methods, however, provide a predictable approach for the generation of gene expression reporter transgenic animals which can be used for quantitative comparisons of gene expression across different animals transfected with the same construct, or for quantitative comparisons of the relative levels of expression of two or more different genes.

10 The present invention solves this and other problems by providing transgenic animals in which at least one single-copy, non-essential gene is replaced with a reporter expression cassette, for example, a luciferase gene operably linked to a heterologous promoter. Thus, the present invention provides novel methods and vector constructs useful for the generation of transgenic animals. The invention further includes methods  
15 of using these animals.

#### SUMMARY OF THE INVENTION

The transgenic animals described herein are useful, for example, when studying *in vivo* regulation of selected genes. Also described herein are methods of generating  
20 populations of substantially isogenic transgenic animals, as well as, vectors useful in these methods.

Accordingly, in one embodiment, the subject invention is directed to a transgenic, non-human mammal, for example, a rodent such as a mouse. The mammal comprises at least one single-copy, non-essential gene in its genome, wherein (i) at least a portion of  
25 at least one single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the gene, and (ii) the polynucleotide sequences comprise a first expression cassette which has been introduced into the mammal or an ancestor of the mammal, at an embryonic stage. The first expression cassette typically comprises a first selectable marker, a first transcriptional promoter element heterologous to the gene, and  
30 light-generating protein coding sequences. The light-generating protein coding sequences are operably linked to the promoter element.

The single-copy, non-essential gene may be selected, for example, from the group consisting of vitronectin, *fosB*, and galactin 3 and the first selectable marker may be selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

In alternative embodiments, the first transcriptional promoter element is an inducible promoter, a repressible promoter, or a constitutive promoter, and may be selected from the group consisting of VEGF, VEGFR, and TIE2.

In additional embodiments, the transgenic, non-human mammal described above comprises a second single-copy, non-essential gene in its genome, wherein (i) at least a portion of the second single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the second gene, and (ii) the polynucleotide sequences comprise a second expression cassette which has been introduced into the mammal or an ancestor of the mammal, at an embryonic stage. The second expression cassette typically comprises a second selectable marker, a second transcriptional promoter element heterologous to the second gene, and light generating protein coding sequences. The light generating protein coding sequences are operably linked to the promoter element.

The first and second transcriptional promoter elements and selectable markers may be the same or different and the light generating protein in the first expression cassette can produce a different color of light relative to the light generating protein in the second expression cassette.

In yet a further embodiment, the invention is directed to a method of producing a transgenic, non-human mammal, such as a mouse. The mammal has at least one single-copy, non-essential gene in its genome. The method comprises

transfecting an embryonic stem cell of the mammal with a linear vector comprising

(a) a first selectable marker and a reporter expression cassette, the reporter expression cassette comprising a transcriptional promoter element operably linked to a light generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, the targeting polynucleotide sequences

flanking (a), wherein (i) the length of the polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) the transcriptional promoter element is heterologous to the single-copy, non-essential gene;

- 5        selecting embryonic stem cells which each have the first selectable marker and reporter expression cassette integrated into its genome;  
              injecting the embryonic stem cells into a host embryo,  
              implanting the embryo in a foster mother,  
              maintaining the foster mother under conditions which allow production of an  
10    offspring that is a transgenic, non-human mammal carrying the reporter expression cassette.

              In certain embodiments, the offspring is capable of germline transmission of the reporter expression cassette and the method may further comprise breeding the offspring with a mammal which is substantially isogenic with the embryonic stem cells, such that  
15    the breeding yields transgenic F1 offspring carrying the reporter cassette. In particular embodiments, the method comprises breeding the first F1 offspring carrying the reporter cassette with a second F1 offspring carrying the reporter cassette, wherein the breeding yields transgenic F2 offspring carrying the reporter cassette.

              In additional embodiments, the embryonic stem cells may be derived from a  
20    mouse having a dark coat color, the mammal substantially isogenic with the embryonic stem cells may have a light coat color, and/or the F2 offspring carrying the reporter cassette may have a light coat color. In particular embodiments, the embryonic stem cells are derived from a C57BL/6 mouse having a dark coat color, and the mammal substantially isogenic with the embryonic stem cells is a C57BL/6-Tyr C2j/+ mouse  
25    having a light coat color.

              In still a further embodiment, the subject invention is directed to a vector for use in generating a transgenic non-human mammal, for example, a rodent such as a mouse. The mammal has at least one single-copy, non-essential gene in its genome. The vector comprises

- 30        (a) a first selectable marker and a reporter expression cassette, the reporter expression cassette comprising a transcriptional promoter element operably linked to a light generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in the mammal's genome, the targeting polynucleotide sequences flanking (a), wherein (i) the length of the targeting polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) the transcriptional promoter element is heterologous to the single-copy, non-essential gene.

In certain embodiments, the first selectable marker provides a positive selection and may be selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase. Additionally, the transcriptional promoter element may be an inducible promoter, a repressible promoter, or a constitutive promoter, and may be selected from the group consisting of VEGF, VEGFR, and TIE2.

In alternative embodiments, the vector further comprises a second selectable marker and at least one target polynucleotide sequence is located between the second selectable marker and the first selectable marker. Additionally, the second selectable marker may provide a negative selection and may be selected from the group consisting of adenosine deaminase, thymidine kinase, and dihydrofolate reductase.

The vectors described above may be circular and may contain at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: target polynucleotide sequence - (a) - targeting polynucleotide sequences or target polynucleotide sequence - (a) - targeting polynucleotide sequences - (second selectable marker).

The coding sequences of the reporter expression cassette present in the vector may comprise codons that are optimal for expression in a host system into which the expression cassette is to be introduced. Additionally, the targeting polynucleotide sequences from single-copy, non-essential genes may be selected from the group consisting of vitronectin, *fosB*, and galactin 3.

The light-generating protein in the mammals and methods described above may be derived from either procaryotic or eucaryotic sources and, in particularly preferred embodiments, the light generating protein is a luciferase.

These and other embodiments of the present invention will be apparent to those of skill in the art in view of the teachings herein.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic depicting construction of the pTK53 vector.

Polynucleotides encoding PGK-P, Neo and TK and 5' and 3' linkers are introduced into  
5 a pKS backbone to produce the vector designated pTK53.

Figure 2 is schematic depicting construction of the pTK-LucR and pTK-LucYG vectors. For pTK-LucR, a polynucleotide encoding LucR is introduced into pTK53. Thus, the pTK-LucR construct contains the PGK-P gene, a neomycin (Neo<sup>r</sup>) gene, a thymidine kinase (TK) gene and sequence encoding red luciferase (Luc-R). For pTK-  
10 LucYG, a polynucleotide encoding LucYG is introduced into pTK53. Thus, the pTK-LucYG construct contains the PGK-P gene, a neomycin (Neo<sup>r</sup>) gene, a thymidine kinase (TK) gene and a sequence encoding yellow-green luciferase (Luc-YG).

Figures 3A is a schematic depicting the vector pTKLR-Vn. Sequences homologous to the vitronectin gene are inserted into pTK-LucR such that they flank the  
15 Neo<sup>r</sup> gene and the Luc-R coding sequence. Figure 3B is a schematic depicting targeting of the linearized pTKLR-Vn vector to the vitronectin chromosomal locus. The VEGF promoter is cloned into the polylinkers between Neo and Luc-R. Upon homologous recombination, the Neo-VEGF-LucR transgene is inserted into the Vn gene. In the figure, (A) shows the targeting vector pTKLR-Vn and (B) shows the mouse vitronectin  
20 gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucR – red luciferase from pGL3Red (Dr. Christopher Contag, Stanford University, Stanford, CA). Regions bearing Vn gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation. Figure 3C shows the  
25 nucleotide sequence of vitronectin (SEQ ID NO:38).

Figure 4A is a schematic depicting the vector pTKLG-Fos. Sequences homologous to the FosB gene are inserted into pTK-LucYG such that they flank the Neo<sup>r</sup> gene and the Luc-YG coding sequence. Figure 4B shows the nucleotide sequence of FosB (SEQ ID NO:39).

30 Figure 5A is a schematic depicting targeting of the linearized pTKLG-Fos vector to the FosB chromosomal locus. The VEGFR2 promoter is cloned into the polylinkers between Neo and Luc-YG. Upon homologous recombination, the Neo-VEGFR2-LucYG



transgene will be inserted into a sequence associated with production of FosB. In the figure, (A) shows the targeting vector, and (B) shows the mouse target gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucYG – yellow green luciferase from pGL3-control vector (Promega, Madison, WI). Regions bearing FosB gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation. Figure 5B is a schematic depicting targeting of the linearized pTKLG-Fos vector to the FosB chromosomal locus. The TIE2 promoter is cloned into the polylinkers between Neo and Luc-R. Upon homologous recombination, the Neo-Tie2-LucYG transgene is inserted into the FosB gene. In the figure, (A) shows the targeting vector, and (B) shows the mouse target gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucYG – yellow green luciferase from pGL3-control vector (Promega). Regions bearing FosB gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation.

Figure 6 depicts PCR conditions for genomic screening for promoters useful in exemplary targeting constructs of the present invention.

Figure 7 depicts generation of targeted transgenic mice using the targeting vectors described herein.

Figure 8 depicts of schematic representation of Southern blot analysis of homologous DNA recombination between pTKLG-Fos targeting vector and the FosB gene.

Figure 9 depicts generation of targeted transgenic mice, using the targeting vectors described herein, and crosses using such transgenics as well as their offspring (F1, first generation; F2, second generation).

Figure 10 depicts crosses using transgenic mice of the present invention to generate dual luciferase transgenic mice.

### MODES FOR CARRYING OUT THE INVENTION

Throughout this application, various publications, patents, and published patent applications are referred to by an identifying citation. The disclosures of these publications, patents, and published patent specifications referenced in this application are included to more fully describe the state of the art to which this invention pertains.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, cell biology and recombinant DNA, which are within the skill of the art. *See, e.g.*, Sambrook, Fritsch, and Maniatis, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd edition (1989); *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, (F.M. Ausubel et al. eds., 1987); the series *METHODS IN ENZYMOLOGY* (Academic Press, Inc.); *PCR 2: A PRACTICAL APPROACH* (M.J. McPherson, B.D. Hames and G.R. Taylor eds., 1995) and *ANIMAL CELL CULTURE* (R.I. Freshney. Ed., 1987).

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise. Thus, for example, reference to "an antigen" includes a mixture of two or more such agents.

#### **Definitions**

As used herein, certain terms will have specific meanings.

The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably to and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown. Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine (T) when the polynucleotide is RNA). Thus, the term polynucleotide sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical

representation can be input into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

A "coding sequence" or a sequence which "encodes" a selected polypeptide, is a  
5 nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide, for example, *in vivo* when placed under the control of appropriate regulatory sequences (or "control elements"). The boundaries of the coding sequence are typically determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but  
10 is not limited to, cDNA from viral, procaryotic or eucaryotic mRNA, genomic DNA sequences from viral or procaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence. Other "control elements" may also be associated with a coding sequence. A DNA sequence encoding a polypeptide can be optimized for expression in a selected cell by using the  
15 codons preferred by the selected cell to represent the DNA copy of the desired polypeptide coding sequence. "Encoded by" refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 to 5 amino acids, more preferably at least 8 to 10 amino acids, and even more preferably at least 15 to 20 amino acids from a  
20 polypeptide encoded by the nucleic acid sequence. Also encompassed are polypeptide sequences which are immunologically identifiable with a polypeptide encoded by the sequence.

Typical "control elements", include, but are not limited to, transcription promoters, transcription enhancer elements, transcription termination signals,  
25 polyadenylation sequences (located 3' to the translation stop codon), sequences for optimization of initiation of translation (located 5' to the coding sequence), translation enhancing sequences, and translation termination sequences. Transcription promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein,  
30 etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters.

"Expression enhancing sequences" typically refer to control elements that improve transcription or translation of a polynucleotide relative to the expression level in the absence of such control elements (for example, promoters, promoter enhancers, enhancer elements, and translational enhancers (e.g., Shine and Delagarno sequences)).

5 "Purified polynucleotide" refers to a polynucleotide of interest or fragment thereof which is essentially free, e.g., contains less than about 50%, preferably less than about 70%, and more preferably less than about 90%, of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example, disruption of the cell  
10 containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

A "heterologous sequence" as used herein is typically refers to either (i) a nucleic acid sequence that is not normally found in the cell or organism of interest, or (ii) a  
15 nucleic acid sequence introduced at a genomic site wherein the nucleic acid sequence does not normally occur in nature at that site. For example, a DNA sequence encoding a polypeptide can be obtained from yeast and introduced into a bacterial cell. In this case the yeast DNA sequence is "heterologous" to the native DNA of the bacterial cell. Alternatively, a promoter sequence from a Tie2 gene can be introduced into the genomic  
20 location of a *fosB* gene. In this case the Tie2 promoter sequence is "heterologous" to the native *fosB* genomic sequence.

A "polypeptide" is used in it broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics. The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used  
25 herein, the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is typically called a polypeptide or a protein.

30 "Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given promoter that is operably linked to a coding sequence (e.g., a reporter expression

cassette) is capable of effecting the expression of the coding sequence when the proper enzymes are present. The promoter or other control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. For example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the coding sequence and the promoter sequence can still be  
5 considered "operably linked" to the coding sequence.

"Recombinant" as used herein to describe a nucleic acid molecule means a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of the polynucleotide  
10 with which it is associated in nature; and/or (2) is linked to a polynucleotide other than that to which it is linked in nature. The term "recombinant" as used with respect to a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. "Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting procaryotic microorganisms or eucaryotic cell  
15 lines cultured as unicellular entities, are used interchangeably, and refer to cells which can be, or have been, used as recipients for recombinant vectors or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement to the original parent, due to  
20 accidental or deliberate mutation. Progeny of the parental cell which are sufficiently similar to the parent to be characterized by the relevant property, such as the presence of a nucleotide sequence encoding a desired peptide, are included in the progeny intended by this definition, and are covered by the above terms. An "isolated polynucleotide" molecule is a nucleic acid molecule separate and discrete from the whole organism with  
25 which the molecule is found in nature; or a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences (as defined below) in association therewith.

Techniques for determining nucleic acid and amino acid "sequence identity" also are known in the art. Typically, such techniques include determining the nucleotide  
30 sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino

acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more sequences (polynucleotide or amino acid) can be compared by determining their "percent identity." The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences  
5 divided by the length of the shorter sequences and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, Atlas of Protein Sequences and Structure, M.O. Dayhoff ed., 5 suppl. 3:353-  
10 358, National Biomedical Research Foundation, Washington, D.C., USA, and normalized by Gribskov, Nucl. Acids Res. 14(6):6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, WI) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis  
15 Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, WI). A preferred method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages  
20 the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is  
25 BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details  
30 of these programs can be found at the following internet address:  
<http://www.ncbi.nlm.gov/cgi-bin/BLAST>. When claiming sequences relative to sequences of the present invention, the desired degrees of sequence identity are at least

80%, 85-90%, preferably 92%, more preferably 95%, and even more preferably 98% sequence identity to the reference sequence (i.e., the sequences of the present invention).

Alternatively, the degree of sequence similarity between polynucleotides can be determined by hybridization of polynucleotides under conditions that form stable  
5 duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when the sequences exhibit at least about 80%-85%, preferably at least about 85%-90%, more preferably at least about 90%-95%, and most preferably at least about 95%-98%  
10 sequence identity over a defined length of the molecules, as determined using the methods above. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that  
15 particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning, supra*; *Nucleic Acid Hybridization, supra*.

Two nucleic acid fragments are considered to "selectively hybridize" as described herein. The degree of sequence identity between two nucleic acid molecules affects the  
20 efficiency and strength of hybridization events between such molecules. A partially identical nucleic acid sequence will at least partially inhibit a completely identical sequence from hybridizing to a target molecule. Inhibition of hybridization of the completely identical sequence can be assessed using hybridization assays that are well known in the art (e.g., Southern blot, Northern blot, solution hybridization, or the like,  
25 see Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.). Such assays can be conducted using varying degrees of selectivity, for example, using conditions varying from low to high stringency. If conditions of low stringency are employed, the absence of non-specific binding can be assessed using a secondary probe that lacks even a partial degree of sequence identity  
30 (for example, a probe having less than about 30% sequence identity with the target molecule), such that, in the absence of non-specific binding events, the secondary probe will not hybridize to the target.

When utilizing a hybridization-based detection system, a nucleic acid probe is chosen that is complementary to a target nucleic acid sequence, and then by selection of appropriate conditions the probe and the target sequence "selectively hybridize," or bind, to each other to form a hybrid molecule. A nucleic acid molecule that is capable of

5 hybridizing selectively to a target sequence under "moderately stringent" typically hybridizes under conditions that allow detection of a target nucleic acid sequence of at least about 10-14 nucleotides in length having at least approximately 70% sequence identity with the sequence of the selected nucleic acid probe. Stringent hybridization conditions typically allow detection of target nucleic acid sequences of at least about 10-

10 14 nucleotides in length having a sequence identity of greater than about 90-95% with the sequence of the selected nucleic acid probe. Hybridization conditions useful for probe/target hybridization where the probe and target have a specific degree of sequence identity, can be determined as is known in the art (see, for example, Nucleic Acid Hybridization: A Practical Approach, editors B.D. Hames and S.J. Higgins, (1985)

15 Oxford; Washington, DC; IRL Press).

With respect to stringency conditions for hybridization, it is well known in the art that numerous equivalent conditions can be employed to establish a particular stringency by varying, for example, the following factors: the length and nature of probe and target sequences, base composition of the various sequences, concentrations of salts and other

20 hybridization solution components, the presence or absence of blocking agents in the hybridization solutions (e.g., formamide, dextran sulfate, and polyethylene glycol), hybridization reaction temperature and time parameters, as well as, varying wash conditions. The selection of a particular set of hybridization conditions is selected following standard methods in the art (see, for example, Sambrook, et al., Molecular

25 Cloning: A Laboratory Manual, Second Edition, (1989) Cold Spring Harbor, N.Y.).

A "vector" is capable of transferring gene sequences to target cells. Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning, and expression

30 vehicles, as well as integrating vectors.

"Nucleic acid expression vector" or "expression cassette" refers to an assembly which is capable of directing the expression of a sequence or gene of interest. The



nucleic acid expression vector includes a promoter which is operably linked to the sequences or gene(s) of interest. Other control elements may be present as well.

Expression cassettes described herein may be contained within a plasmid construct. In addition to the components of the expression cassette, the plasmid construct may also  
5 include a bacterial origin of replication, one or more selectable markers, a signal which allows the plasmid construct to exist as single-stranded DNA (e.g., a M13 origin of replication), a multiple cloning site, and a "mammalian" origin of replication (e.g., a SV40 or adenovirus origin of replication).

An "expression cassette" comprises any nucleic acid construct capable of  
10 directing the expression of a gene/coding sequence of interest. Such cassettes can be constructed into a "vector," "vector construct," "expression vector," or "gene transfer vector," in order to transfer the expression cassette into target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

"Luciferase," unless stated otherwise, includes prokaryotic and eukaryotic  
15 luciferases, as well as variants possessing varied or altered optical properties, such as luciferases that produce different colors of light (e.g., Kajiyama, N., and Nakano, E., *Protein Engineering* 4(6):691-693 (1991)).

"Light-generating" is defined as capable of generating light through a chemical reaction or through the absorption of radiation.

20 A "light generating protein" or "light-emitting protein" is a protein capable of generating light in the visible spectrum (between approximately 350 nm and 800 nm). Examples include bioluminescent proteins such as luciferases, e.g., bacterial and firefly luciferases, as well as fluorescent proteins such as green fluorescent protein (GFP).

"Light" is defined herein, unless stated otherwise, as electromagnetic radiation  
25 having a wavelength of between about 300 nm and about 1100 nm.

"Animal" as used herein typically refers to a non-human mammal, including, without limitation, farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens,  
30 turkeys and other gallinaceous birds, ducks, geese, and the like. The term does not denote a particular age. Thus, both adult and newborn individuals are intended to be covered.

A "transgenic animal" refers to a genetically engineered animal or offspring of genetically engineered animals. A transgenic animal usually contains material from at least one unrelated organism, such as from a virus, plant, or other animal. The "non-human animals" of the invention include vertebrates such as rodents, non-human  
5 primates, sheep, dogs, cows, amphibians, birds, fish, insects, reptiles, etc. The term "chimeric animal" is used to refer to animals in which the heterologous gene is found, or in which the heterologous gene is expressed in some but not all cells of the animal.

"Analyte" as used herein refers to any compound or substance whose effects (*e.g.*, induction or repression of a specific promoter) can be evaluated using the test  
10 animals and methods of the present invention. Such analytes include, but are not limited to, chemical compounds, pharmaceutical compounds, polypeptides, peptides, polynucleotides, and polynucleotide analogs. Many organizations (*e.g.*, the National Institutes of Health, pharmaceutical and chemical corporations) have large libraries of chemical or biological compounds from natural or synthetic processes, or fermentation  
15 broths or extracts. Such compounds/analytes can be employed in the practice of the present invention.

As used herein, the term "positive selection marker" refers to a gene encoding a product that enables only the cells that carry the gene to survive and/or grow under certain conditions. For example, plant and animal cells that express the introduced  
20 neomycin resistance (Neo<sup>r</sup>) gene are resistant to the compound G418. Cells that do not carry the Neo<sup>r</sup> gene marker are killed by G418. Other positive selection markers will be known to those of skill in the art. Typically, positive selection markers encode products that can be readily assayed. Thus, positive selection markers can be used to determine whether a particular DNA construct has been introduced into a cell, organ or tissue.

25 "Negative selection marker" refers to gene encoding a product which can be used to selectively kill and/or inhibit growth of cells under certain conditions. Non-limiting examples of negative selection inserts include a herpes simplex virus (HSV)-thymidine kinase (TK) gene. Cells containing an active HSV-TK gene are incapable of growing in the presence of gangcylovir or similar agents. Thus, depending on the substrate, some  
30 gene products can act as either positive or negative selection markers.

The term "homologous recombination" refers to the exchange of DNA fragments between two DNA molecules or chromatids at the site of essentially identical nucleotide

sequences. It is understood that substantially homologous sequences can accommodate insertions, deletions, and substitutions in the nucleotide sequence. Thus, linear sequences of nucleotides can be essentially identical even if some of the nucleotide residues do not precisely correspond or align (see, above).

5           A "knock-out" mutation refers to partial or complete loss of expression of at least a portion the target gene. Examples of knock-out mutations include, but are not limited to, gene-replacement by heterologous sequences, gene disruption by heterologous sequences, and deletion of essential elements of the gene (e.g., promoter region, portions of a coding sequence). A "knock-out" mutation is typically identified by the phenotype  
10           generated by the mutation.

          A "single-copy gene" as used herein refers to a gene represented in an organism's genome only by a single copy at a particular chromosomal locus. Accordingly, a diploid organism has two copies of the gene and both copies occur at the same chromosomal location.

15           A "gene" as used in the context of the present invention is a sequence of nucleotides in a genetic nucleic acid (chromosome, plasmid, etc.) with which a genetic function is associated. A gene is a hereditary unit, for example of an organism, comprising a polynucleotide sequence (e.g., a DNA sequence for mammals) that occupies a specific physical location (a "gene locus" or "genetic locus") within the  
20           genome of an organism. A gene can encode an expressed product, such as a polypeptide or a polynucleotide (e.g., tRNA). Alternatively, a gene may define a genomic location for a particular event/function, such as the binding of proteins and/or nucleic acids (e.g., phage attachment sites), wherein the gene does not encode an expressed product. Typically, a gene includes coding sequences, such as, polypeptide encoding sequences,  
25           and non-coding sequences, such as, promoter sequences, poly-adenylation sequences, transcriptional regulatory sequences (e.g., enhancer sequences). Many eucaryotic genes have "exons" (coding sequences) interrupted by "introns" (non-coding sequences). In certain cases, a gene may share sequences with another gene(s) (e.g., overlapping genes).

          "Isogenic" means two or more organisms or cells that are considered to be  
30           genetically identical. "Substantially isogenic" means two or more organisms or cells wherein, at the majority of genetic loci (e.g., greater than 99.000%, preferably more than 99.900%, more preferably greater than 99.990%, even more preferably greater than

99.999%), there exists genetic identity between the organisms or cells being compared. In the context of the present invention, two organisms (for example, mice) are considered to be "substantially isogenic" if, for example, inserted transgenes are the primary differences between the genetic make-up of the mice being compared. Further, if, for example, the genetic backgrounds of the mice being compared are the same with the exception that one of the mice has one or several defined mutation(s) (for example, affecting coat color), then these mice are considered to be substantially isogenic. An example of two strains of substantially isogenic mice are C57BL/6 and C57BL/6-Tyr C2j/+.

10 A "pseudogene" as used herein, refers to a type of gene sequence found in the genomes, typically, of eucaryotes, where the sequence closely resembles a known functional gene, but differs in that the pseudogene is non-functional. For example, the pseudogene sequence may contain several stop codons in what would correspond to an open reading frame in the functional gene. Pseudogenes can also have deletions or  
15 insertions relative to their corresponding functional gene. If, for example, in a genome there is a functional gene and a related pseudogene, the functional gene is considered to be a single-copy gene (accordingly, the pseudogene is considered to be single-copy as well).

A "non-essential gene" refers to a gene whose deletion, disruption, elimination,  
20 reduction of gene function, or mutation is non-lethal, and does not obviously adversely affect the organisms' ability to mature and reproduce. A "non-essential gene with no phenotype" refers to a non-essential gene whose deletion, disruption, elimination, reduction of gene function or mutation has no deleterious effect on the organism. Typically there are no phenotypically reflected gene dosage effects associated with  
25 modification of a non-essential gene with no phenotype -- for example, deletion, disruption or mutation of both copies of a non-essential gene with no phenotype in a diploid organism has essentially the same effect as deletion, disruption, or mutation of one of the two copies present in the diploid organism. In the context of the present invention, a non-essential gene is typically one whose function has been eliminated (e.g.,  
30 by a deletion mutation) and such elimination of function was non-lethal and the organism developed, matured, and was able to reproduce.

The "native sequence" or "wild-type sequence" of a gene is the polynucleotide sequence that comprises the genetic locus corresponding to the gene, e.g., all regulatory and open-reading frame coding sequences required for expression of a completely functional gene product as they are present in the wild-type genome of an organism. The native sequence of a gene can include, for example, transcriptional promoter sequences, translation enhancing sequences, introns, exons, and poly-A processing signal sites. It is noted that in the general population, wild-type genes may include multiple prevalent versions that contain alterations in sequence relative to each other and yet do not cause a discernible pathological effect. These variations are designated "polymorphisms" or "allelic variations."

By "replacement sequence" is meant a polynucleotide sequence that is substituted for at least a portion of the native or wild-type sequence of a gene.

"Linear vector" or "linearized vector," as used herein, is a vector having two ends. For example, circular vectors, such as plasmids, can be linearized by digestion with a restriction endonuclease that cuts at a single site in the plasmid. Preferably, the targeting vectors described herein are linearized such that the ends are not within the targeting sequences.

### General Overview

The present invention relates to vector constructs and methods of creating transgenic animals to be used, for example, as test systems. Methods of using the animals of the present invention include, but are not limited to, using these animals for studies involving tumor growth and other disease conditions. In the practice of the present invention, transgenic, non-human mammals are constructed where a single-copy, non-essential gene is replaced by a reporter expression cassette, preferably a gene encoding a light-generating protein, such as a luciferase-encoding gene, operably linked to a promoter. A variety of promoters are useful in the practice of the present invention, for example, promoters derived from genes associated with tumorigenesis or angiogenesis. Thus, an exemplary promoter can be one that is associated with proteins induced during tumorigenesis, for instance in the presence of tumor generating compounds or of tumors themselves. In this way, expression of the reporter cassette is induced in the animal when, for example, tumors are present, and progression of the

tumor can be evaluated by non-invasive imaging methods using the whole animal.

Another exemplary promoter is one that is derived from a gene associated with angiogenesis. Because the promoter is linked to a reporter such as luciferase, non-invasive monitoring of the progression of angiogenesis is possible.

5           Various forms of the different embodiments of the invention, described herein, may be combined.

Non-invasive imaging and/or detecting of light-emitting conjugates in mammalian subjects was described in U.S. Patent No. 5,650,135, by Contag, et al., issued 22 July 1997. This imaging technology can be used in the practice of the present  
10   invention in view of the teachings of the present specification. In the imaging method, the conjugates contain a biocompatible entity and a light-generating moiety. Biocompatible entities include, but are not limited to, small molecules such as cyclic organic molecules; macromolecules such as proteins; microorganisms such as viruses, bacteria, yeast and fungi; eukaryotic cells; all types of pathogens and pathogenic  
15   substances; and particles such as beads and liposomes. In another aspect, biocompatible entities may be all or some of the cells that constitute the mammalian subject being imaged, for example, cells carrying the vector constructs of the present invention expressing a reporter expression cassette.

Light-emitting capability is conferred on the biocompatible entities by the  
20   conjugation of a light-generating moiety. Such moieties include fluorescent molecules, fluorescent proteins, enzymatic reactions giving off photons and luminescent substances, such as bioluminescent proteins. In the context of the present invention, light emitting capability is typically conferred on target cells by having at least one copy of a light-generating protein, *e.g.*, a luciferase, present. In preferred embodiments, luciferase is  
25   operably linked to appropriate control elements which can facilitate expression of a polypeptide having luciferase activity. Substrates of luciferase can be endogenous to the cell or applied to the cell or system (*e.g.*, injection into a transgenic mouse, having cells carrying a luciferase construct, of a suitable substrate for the luciferase, for example, luciferin). The conjugation may involve a chemical coupling step, genetic engineering of  
30   a fusion protein, or the transformation of a cell, microorganism or animal to express a light-generating protein.

### Targeting Constructs

The targeting cassettes described herein typically include the following components: (1) a suitable vector backbone; (2) a polynucleotide encoding a light generating protein (3) a promoter operably linked to the light generating protein-  
5 encoding gene, wherein the promoter is heterologous to the light generating protein coding sequences; (4) a sequence encoding a positive selection marker; (5) insertion sites flanking the sequence encoding the positive selection marker and the polynucleotide encoding a light generating protein gene, for insertion of sequences which target a single-copy, non-essential chromosomal gene; and, optionally, (6) a sequence encoding a  
10 negative selection marker. Exemplary targeting constructs are shown in Figures 3B, 5A and 5B and described in Examples 1-3.

Suitable vector backbones generally include an F1 origin of replication; a colE1 plasmid-derived origin of replication; polyadenylation sequence(s); sequences encoding antibiotic resistance (*e.g.*, ampicillin resistance) and other regulatory or control elements.  
15 Non-limiting examples of appropriate backbones include: pBluescriptSK (Stratagene, La Jolla, CA); pBluescriptKS (Stratagene, La Jolla, CA) and other commercially available vectors.

In one aspect of the invention the light generating protein is luciferase. Luciferase coding sequences useful in the practice of the present invention include, but  
20 are not limited to, sequences obtained from *lux* genes (procaryotic genes encoding a luciferase activity) and *luc* genes (eucaryotic genes encoding a luciferase activity). A variety of luciferase encoding genes have been identified including, but not limited to, the following: B.A. Sherf and K.V. Wood, U.S. Patent No. 5,670,356, issued 23 September 1997; Kazami, J., et al., U.S. Patent No. 5,604,123, issued 18 February 1997;  
25 S. Zenno, et al, U.S. Patent No. 5,618,722; K.V. Wood, U.S. Patent No. 5,650,289, issued 22 July 1997; K.V. Wood, U.S. Patent No. 5,641,641, issued 24 June 1997; N. Kajiyama and E. Nakano, U.S. Patent No. 5,229,285, issued 20 July 1993; M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,292,658, issued 8 March 1994; M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,418,155, issued 23 May 1995; de Wet, J.R., et al,  
30 *Molec. Cell. Biol.* 7:725-737, 1987; Tatsumi, H.N., et al, *Biochim. Biophys. Acta* 1131:161-165, 1992; and Wood, K.V., et al, *Science* 244:700-702, 1989. Eukaryotic luciferase catalyzes a reaction using luciferin as a luminescent substrate to produce light,

whereas prokaryotic luciferase catalyzes a reaction using an aldehyde as a luminescent substrate to produce light.

Wild-type firefly luciferases typically have an emission maxima at about 550 nm. Numerous variants with differing emission maxima have also been studied. For example, Kajiyama and Nakano (*Protein Eng.* 4(6):691-693, 1991; U.S. Patent No. 5,330,906, issued 19 July 1994) teach five variant firefly luciferases generated by single amino acid changes to the *Luciola cruciata* luciferase coding sequence. The variants have emission peaks of 558 nm, 595 nm, 607 nm, 609 nm and 612 nm. A yellow-green luciferase with an emission peak of about 540 nm is commercially available from Promega, Madison, WI under the name pGL3. A red luciferase with an emission peak of about 610 nm is described, for example, in Contag et al. (1998) *Nat. Med.* 4:245-247 and Kajiyama et al. (1991) *Prot. Eng.* 4:691-693.

Positive selection markers include any gene which a product that can be readily assayed. Examples include, but are not limited to, a hprt gene (Littlefield, J. W., *Science* 145:709-710 (1964)), a xanthine-guanine phosphoribosyltransferase (gpt) gene, or an adenosine phosphoribosyltransferase (aprt) gene (Sambrook et al., *supra*), a thymidine kinase gene (i.e. "TK") and especially the TK gene of herpes simplex virus (Giphart-Gassler, M. et al., *Mutat. Res.* 214:223-232 (1989)), a nptII gene (Thomas, K. R. et al., *Cell* 51:503-512 (1987); Mansour, S. L. et al., *Nature* 336:348-352 (1988)), or other genes which confer resistance to amino acid or nucleoside analogues, or antibiotics, etc, for example, gene sequences which encode enzymes such as dihydrofolate reductase (DHFR) enzyme, adenosine deaminase (ADA), asparagine synthetase (AS), hygromycin B phosphotransferase, or a CAD enzyme (carbamyl phosphate synthetase, aspartate transcarbamylase, and dihydroorotase). Addition of the appropriate substrate of the positive selection marker can be used to determine if the product of the positive selection marker is expressed, for example cells which do not express the positive selection marker nptII, are killed when exposed to the substrate G418 (Gibco BRL Life Technology, Gaithersburg, MD).

The targeting vector typically contains insertion sites for inserting targeting sequences (e.g., sequences that are substantially homologous to the target sequences in the host genome where integration of the targeting vector/expression cassette is desired). These insertion sites are preferably included such that there are two sites, one site on



either side of the sequences encoding the positive selection marker, light generating protein (e.g., luciferase) and the promoter. Insertion sites are, for example, restriction endonuclease recognition sites, and can, for example, represent unique restriction sites. In this way, the vector can be digested with the appropriate enzymes and the targeting  
5 sequences ligated into the vector.

Optionally, the targeting construct can contain a polynucleotide encoding a negative selection marker. Suitable negative selection markers include, but are not limited to, HSV-tk (see, e.g., Majzoub et al. (1996) *New Engl. J. Med.* 334:904-907 and U.S. Patent No. 5,464,764), as well as genes encoding various toxins including the  
10 diphtheria toxin, the tetanus toxin, the cholera toxin and the pertussis toxin. A further negative selection marker gene is the hypoxanthine-guanine phosphoribosyl transferase (HPRT) gene for negative selection in 6-thioguanine.

Exemplary promoters and single-copy, non-essential genes for use in the vector constructs and methods of the present invention are described below.

15

### Promoters

The targeting constructs and transgenic animals described herein contain a sequence encoding a light generating protein, e.g., luciferase, gene operably linked to a promoter. The promoter may be from the same species as the transgenic animal (e.g.,  
20 mouse promoter used in construct to make transgenic mouse) or from a different species (e.g., human promoter used in construct to make transgenic mouse). The promoter can be derived from any gene of interest. In one embodiment of the present invention, the promoter is derived from a gene whose expression is induced during angiogenesis, for example pathogenic angiogenesis like tumor development. Thus, when a tumor begins  
25 to develop in a transgenic animal carrying a vector construct of the present invention, the promoter is induced and the animal expresses luciferase, which can then be monitored *in vivo*.

Exemplary promoters for use in the present invention are selected such that they are functional in a cell type and/or animal into which they are being introduced.

30 Exemplary promoters include, but are not limited to, promoters obtained from the following mouse genes: vascular endothelial growth factor (VEGF) (VEGF promoter described in U.S. Patent No. 5,916,763; Shima et al. (1996) *J. Bio. Chem.* 271:3877-

3883; sequence available on NCBI under accession number U41383); VEGFR2, also known as Flk-1, (VEGFR-2 promoter described, for example, in Röncke et al. (1996) *Circ. Res.* 79:277-285; Patterson et al. (1995) *J. Bio. Chem.* 270:23111-23118; Kappel et al. (1999) *Blood* 93:4282-4292; sequence available as accession number X89777 of NCBI database); Tie2, also known as Tek (Tie2 promoter described, for example, in Fadel et al. (1998) *Biochem. J.* 338:335-343; Schlaeger et al. (1995) *Develop.* 121:1089-1098; Schlager et al. (1997) *PNAS USA* 94:3058-3063). VEGF is a specific mitogen for EC *in vitro* and a potent angiogenic factor *in vivo*. In a tumorigenesis study, it was shown that VEGF was critical for the initial subcutaneous growth of T-47D breast carcinoma cells transplanted into nude mice, whereas other angiogenic factors, such as, bFGF can compensate for the loss of VEGF after the tumors have reached a certain size (Yoshiji, H., et al., 1997 *Cancer Research* 57: 3924-28). VEGF is a major mediator of aberrant EC proliferation and vascular permeability in a variety of human pathologic situation, such as, tumor angiogenesis, diabetic retinopathy and rheumatoid arthritis (Benjamin LE, et al., 1997 *PNAS* 94: 8761-66; Soker, S., et al., 1998 *Cell* 92: 735-745). VEGF is synthesized by tumor cells *in vivo* and accumulates in nearby blood vessels. Because leaky tumor vessels initiate a cascade of events, which include plasma extravasation and which lead ultimately to angiogenesis and tumor stroma formation, VEGF plays a pivotal role in promoting tumor growth (Dvorak, H.F., et al., 1991 *J Exp Med* 174:1275-8). VEGF expression was upregulated by hypoxia (Shweiki, D., et al., 1992 *Nature* 359: 843-5). VEGF is also upregulated by overexpression of v-Src oncogene (Mukhopadhyay, D., et al., 1995 *Cancer Res.* 15: 6161-5), c-SRC (Mukhopadhyay, D., et al., 1995 *Nature* 375: 577-81), and mutant ras oncogene (Plate, K.H., et al., 1992 *Nature* 359: 845-8). The tumor suppressor p53 downregulates VEGF expression (Mukhopadhyay, D., et al., 1995 *Cancer Res.* 15: 6161-5).

A number of cytokines and growth factors, including PGF and TPA (Grugel, S., et al., 1995 *J. Biological Chem.* 270: 25915-9), EGF, TGF- $\beta$ , IL-1, IL-6 induce VEGF mRNA expression in certain type of cells (Ferrara, N., et al., 1997 *Endocr. Rev.* 18: 4-25). Kaposi's sarcoma-associated herpesvirus (KSHV) encoded a G-protein-coupled receptor, a homolog of IL-8 receptor, can activate JNK/SAPK and p38MAPK and increase VEGF production, thus causing cell transformation and tumorigenicity (Bais, C., et al., *Nature* 1998 391:86-9). VEGF overexpression in skin of transgenic mice

induces angiogenesis, vascular hyperpermeability and accelerated tumor development (Larcher, F., et al., *Oncogene* 1998 17:303-11).

VEGF-B (cDNA sequences available on databases) is a mitogen for EC and may be involved in angiogenesis in muscle and heart (Olofsson, B., et al., 1996 *Proc Natl Acad Sci U S A* 93:2576-81). Shown *in vitro*, binding of VEGF-B to its receptor VEGFR-1 leads to increased expression and activity of urokinase type plasminogen activator and plasminogen activator inhibitor, suggesting a role for VEGF-B in the regulation of extracellular matrix degradation, cell adhesion, and migration (Olofsson, B., et al., 1998 *Proc Natl Acad Sci U S A* 95:11709-14).

VEGF-C (see, e.g., U.S. Patent No. 5,916,763 and Shima et al., *supra*) may regulate angiogenesis of lymphatic vasculature, as suggested by the pattern of VEGF-C expression in mouse embryos (Kukk, E., et al., 1996 *Development* 122: 3829-37). Although VEGF-C is also a ligand for VEGFR-2, the functional significance of this potential interaction is unknown. Overexpression of VEGF-C in the skin of transgenic mice resulted in lymphatic, but not vascular, endothelial proliferation and vessel enlargement, suggesting the major function of VEGF-C is through VEGFR-3 rather than VEGFR-2 (Jeltsch M, et al., 1997 *Science* 276:1423-5). Shown by the CAM assay, VEGF and VEGF-C are specific angiogenic and lymphangiogenic growth factors, respectively (Oh, S.J., et al., (1997) *Devel. Biol.* 188: 96-109). VEGF-C overexpression in the skin of transgenic mice resulted in lymphatic, but not vascular, endothelial proliferation and vessel enlargement (Jeltsch M, et al., 1997 *Science* 276:1423-5).

VEGF-D (cDNA sequences available on databases) is a mitogen for EC. Given that VEGF-D can also activate VEGFR-3, it is possible that VEGF-D could be involved in the regulation of growth and/or differentiation of lymphatic endothelium (Achen, M.G., et al., 1998 *Proc Natl Acad Sci U S A* 95: 548-53). VEGF-D is induced by transcription factor c-Fos in mouse (Orlandini, M., 1996 *PNAS* 93: 11675-80).

VEGFR-1 signaling pathway may regulate normal endothelial cell-cell or cell matrix interactions during vascular development, as suggested by the knockout study (Fong, G.H., et al., 1995 *Nature* 376: 65-69). Although VEGFR-1 has a higher affinity to VEGF than VEGFR-2, it does not transduce the mitogenic signals of VEGF in ECs (Soker, S., et al., 1998 *Cell* 92: 735-745). VEGFR-2 (see, e.g., Röncke et al., Patterson et al., Kappel et al. (1999), *supra*) appears to be the major transducer of VEGF signals in

EC that result in chemotaxis, mitogenicity and gross morphological changes in target cells (Soker, S., et al., 1998 *Cell* 92: 735-745). VEGFR-3 has an essential role in the development of the embryonic cardiovascular system before the emergence of lymphatic vessels, as shown by the knockout study (Dumont, D.J., et al., 1998 *Science* 282: 946-949). Neuropilin-1 (see, e.g., Soker et al. (1998) *Cell* 92:735-745) is a receptor for VEGF165. It can enhance the binding of VEGF165 to VEGFR-2 and VEGF165 mediated chemotaxis (Soker, S., et al., 1998 *Cell* 92: 735-745). Neuropilin1 overexpression in transgenic mice resulted in embryonic lethality. The embryos possessed excess capillaries and blood vessels. Dilated vessels and hemorrhage were also observed (Kitsukawa, T., et al., 1995 *Development* 121: 4309-18).

Further promoters of interest include, but are not limited to, the following. Ang2 is expressed only at predominant vascular remodeling sites, such as ovary, placenta, uterus (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). In glioblastoma angiogenesis, Ang2 is found to be expressed in endothelial cells of small blood vessel and capillaries while Ang1 is expressed in glioblastoma tumor cells (Stratmann, A., 1998 *Am J Pathol* 153: 1459-66). Ang2 is up-regulated in bovine microvascular endothelial by VEGF, bFGF, cytokines, hypoxia (Mandriota, S.J., 1998 *Circ Res* 83: 852-9). Ang2 transgenic overexpression disrupts angiogenesis, and is embryonic lethal (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). Ang1 is widely expressed, less abundant in heart and liver (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). Ang1 is expressed in mesenchymal cells and may up-regulate the expression of Tie2 in the endothelial cells (Suri, C., et al., 1996 *Cell* 87: 1171-1180). Ang1 overexpression in the skin of transgenic mice produces larger, more numerous, and more highly branched vessels (Suri, C., et al., *Science* 1998 282:468-71). Tie2 (see, e.g., Fadel et al.; Schlaeger et al. (1995), and Schlaeger et al. (1997), *supra*) is endothelial cell specific, up-regulated during wound healing, follicle maturation (Puri, M.C., et al., 1995 *EMBO J* 14: 5884-91) and pathologic angiogenesis (Kaipainen, A., 1994 *Cancer Research* 54: 6571-77), such as, glioblastoma (Stratmann, A., 1998 *Am J Pathol* 153: 1459-66). Tie2 is also expressed in non-proliferating adult endothelium and endothelial cell lines (Dumont, D.J., et al. (1994) *Genes & Develop.* 8:1897-1909). A Tie2 activating mutation causes vascular dysmorphogenesis (Vikkula M, et al., 1996 *Cell* 87: 1181-1190). Tie2 mutant overexpression in transgenic mice is embryonic lethal (Dumont, D.J., et al., *supra*).

Other promoters useful in the practice of the present invention include, by way of example, promoters derived from the sequences encoding the following polypeptide products: PTEN (dual specificity phosphatase); BAI (brain-specific angiogenesis inhibitor); KAI1 (KANGAI 1); catenin beta-1 (cadherin-associated protein, beta); COX2 (PTGS2 cyclooxygenase 2, a.k.a. prostaglandin-endoperoxide synthase 2); MMP2 (72 kDa Type IV-A collagenase); MMP9 (92 kDa type IV-B collagenase); TIMP2 (tissue inhibitor of metalloproteinase 2); and TIMP3 (tissue inhibitor of metalloproteinase 3).

PTEN is a tumor suppressor gene and encodes a protein of 403 amino acids. (Li et al. (1997) *Science* 275:1943-1946; DiCristofano et al. (1998) *Nature Genet.* 19:348-355). Overexpression of PTEN has been shown to inhibit cell migration and it is postulated that this protein may function as a tumor suppressor by negatively regulating cell interactions with the extracellular matrix or by negatively regulating the PI3K/PKB/Akt signaling pathway. (Tamura et al. (1998) *Science* 280:1614-1617; Stambolic et al. (1998) *Cell* 95:29-29). Mutations in PTEN have been detected in cancer cell lines and in the germline of patients having Cowden disease, Lhermitte-Duclos disease and Bannayan-Zonana syndrome (diseases and syndromes which are characterized by hyperplastic/dysplastic changes in the prostate, skin and colon and which are associated with an increased risk of certain cancers, for example, breast cancer, prostate cancer and colon cancer). (Marsh et al. (1998) *Hum. Molec. Genet.* 7:507-515; Marsh et al. (1998) *J. Med. Genet.* 35:881-885; Nelen et al. (1997) *Hum. Molec. Genet.* 6:1383-1387).

BAI1 protein is predicted to be 1,584 amino acids in length and includes an extracellular domain, an intracellular domain and a 7-span transmembrane region similar to that of the secretin receptor. (Nishimori et al. (1997) *Oncogene* 15:245-2150). The extracellular region of BAI1 has a single Arg-Gly-Asp (RGD) motif recognized by integrins and also has five sequences corresponding to the thrombospondin type I (accession number 188060) repeats that can inhibit angiogenesis includes by basic fibroblast growth factor (bFGF, accession number 134920). Shiratsuchi et al. (1997) *Cytogenet. Cell Genet.* 79:103-108, cloned 2 other brain-specific angiogenesis inhibiting genes, designated BAI2 (accession number 602683) and BAI 3 (accession number 602684). Thus, it is postulated that members of this gene family may play a role in suppression of glioblastoma.

KAI1 encodes a 267 amino acid protein which is a member of the leukocyte surface glycoprotein family. The protein has 4 hydrophobic transmembrane domains and 1 large extracellular hydrophilic domain with three potential N-glycosylation sites. (Dong et al. (1995) *Science* 268:884-886). Molecular analysis of KAI1 is described, for example, in Dong et al. (1997) *Genomics* 41:25-32. KAI1 is a tumor metastasis suppressor gene that is capable of inhibiting the metastatic process in experimental animals. Expression of KAI1 is downregulated during tumor progression of prostate, breast, lung, bladder and pancreatic cancers in humans, apparently at the transcriptional or postranscriptional level. Mashimo et al. (1998) *PNAS USA* 95:11307-11311, found that the tumor suppressor gene p53 can directly inactivate the KAI1 gene by interacting with the region 5' to the coding sequence, suggesting a direct relationship between p53 and KAI1.

Catenin beta-1 is an adherens junction (AJ) protein, which are critical for establishing and maintaining epithelial cell layers, for instance during embryogenesis, wound healing and tumor cell metastasis. Molecular analysis, including description of sequence homology to plakoglobin (accession number 173325), homology to the drosophila gene "armadillo" and interactions with Lef1/Tcf DNA binding proteins, is described, for example, in Nollet et al. (1996) *Genomics* 32:413-424; McCrea et al. (1991) *Science* 254:1359-1361 and Korinek et al. (1997) *Science* 275:1784-1787. In addition, studies by Korinek et al., *supra* and Morin et al. (1997) *Science* 275:1787-1790, have indicated that APC (accession number 175100) negatively regulates catenin beta and that regulation of this protein is critical to the tumor suppressive effect of APC. Abnormally high levels of beta-catenin have been detected in certain human melanoma cell lines. (Rubinfeld et al. (1997) *Science* 275:1790-1792. Koch et al. (1999) *Cancer Res.* 59:269-273 report that childhood hepatoblastomas frequently carry a mutated degradation targeting box of the beta-catenin gene. Transgenic mice which express catenin beta under the control of an epidermal promoter undergo de novo hair morphogenesis and eventually these animals develop two types of tumors -- epithelioid cysts and trichofolliculomas. Gat et al. (1998) *Cell* 95:605-614.

COX2 encodes a cyclooxygenase and is a key regulator of prostaglandin synthesis. (Hla et al. (1992) *PNAS USA* 89:7384-7388; Jones et al. (1993) *J. Biol. Chem.* 268:9049-9054). In particular, COX2 is generally considered to be a mediator of

inflammation and overexpression of COX2 in rat epithelial cells results in elevated levels of E-cadherin and Bcl2. (Tsuji & DuBois (1995) *Cell* 83:493-501). In co-cultures of endothelial cells and colon carcinoma cells, cells that overexpress COX2 produce prostaglandins, proangiogenic factors and stimulate both endothelial migration and tube formation. (Tsuji et al. (1998) *Cell* 93:705-716). Experiments conducted using APC knock-out mice have demonstrated that animals homozygous for a disrupted COX2 locus develop significantly more adenomatous polyps. (Oshima et al. (1996) *Cell* 87:803-809). COX-2 "knock out" mice develop severe nephropathy, are susceptible to peritonitis, exhibit reduced arachidonic acid-induced inflammation and exhibit reduced indomethacin-induced gastric ulceration. (Morham et al. (1995) *Cell* 83:473-482; Langenbach et al. (1995) *Cell* 83:483-492). Female mice that are deficient in cyclooxygenase 2 exhibit multiple reproductive failures. (Lim et al. (1997) *Cell* 91:197-208).

MMP2 is a metalloproteinase that specifically cleaves type IV collagen. A C-terminal fragment of MMP2, termed PEX, prevents normal binding to alpha-V/beta-3 and disrupts angiogenesis and tumor growth. (Brooks et al. (1998) *Cell* 92:391-400).

MMP9 is a collagenase secreted from normal skin fibroblasts. MMP9 null mice exhibit an abnormal pattern of skeletal growth plate vascularization and ossification. (Vu et al. (1998) *Cell* 93:411-422).

TIMP2 is a collagenase and appears to play a major role in modulating the activity of interstitial collagenase and a number of connective tissue metalloendoproteases. (Stetler-Stevenson et al. (1989) *J. Biol. Chem.* 264:17372-17378). Unlike TIMP1 and TIMP3, TIMP2 is not upregulated by TPA or TGF-beta. (Hammani et al. (1996) *J. Biol. Chem.* 271:25498-25505).

TIMP3 (Wilde et al. (1994) *DNA Cell Biol.* 13:711-718) is localized in the extracellular matrix in both its glycosylated and unglycosylated forms. Studies of mutant TIMP3 proteins have demonstrated that C-terminal truncations do not bind to the extracellular matrix. (Langton et al. (1998) *J. Biol. Chem.* 273:16778-16781).

As one of skill in the art will appreciate in view of the teachings of the present specification, promoter sequences can be easily derived and isolated from known polypeptide sequences or from cDNA or genomic sequences, using method known in the art in view of the teachings herein. An exemplary method of isolating promoter

sequences using cDNA is via a GenomeWalker® kit, commercially available from Clontech (Palo Alto, CA), and described on page 27 of the 1997-1998 Clontech catalog.

#### Targeting Sequences: Non-Essential Genes

5 Central to the present invention is the fact that the targeting constructs contain "targeting" sequences (flanking, for example, the light generating protein-encoding sequence and promoter) derived from a single-copy, non-essential gene. These targeting sequences in the construct act via homologous recombination to replace at least a portion of the non-essential gene in the genome with the light-generating protein-encoding (*e.g.*,  
10 luciferase-encoding) sequence operably linked to a promoter.

Non-limiting examples of targeting sequences for use in generating transgenic mice include sequences obtained from or derived from vitronectin, Fos B and galactin 3. A search of Mouse Knockout & Mutation Database (Genome Systems, Inc., St. Louis, MO) can be used to identify genes that have been knocked-out in mice where the  
15 generated knockout mice displayed no obvious defects. The chromosomal locus for all these genes can be used to target promoter-(light generating protein, *e.g.*, luciferase) transgenes similar to what is described in Example 2. Single-copy, non-essential mouse genes identified in this manner include, but are not limited to, the following: Moesin (Msn), Doi Y., et al., J Biol Chem 1999, 274:2315-2321; Plasminogen activator  
20 inhibitor, type II (Planh2) and Planh1, Dougherty K.M., Proc Natl Acad Sci USA 1999, 96:686-691; Protein tyrosine phosphatase, receptor type, B (Ptpnb), Elchebly et al. (1999) Science 283:1544-1548; Presenilin 1 (Psen1), Guo Q, et al. (1999) Proc Natl Acad Sci USA, 96:4125-4130; Protein kinase, mitogen-activated 9 (Prkm9) / SAPK/Erk/kinase 2 (Serk2), Kuan CY et al. (1999) Neuron 4:667-676; CD152 antigen  
25 (Cd152) / CD86 antigen (Cd86) / CD80 antigen (Cd80), Mandelbrot DA, et al. (1999) J Exp Med, 189:435-440; Poly (ADP-ribose) polymerase (Adprp), Masutani M, et al. (1999), Proc Natl Acad Sci USA 96:2301-2304; Sodium channel, nonvoltage-gated 1 beta (Scnn1b), Pradervand S, et al. (1999) Proc Natl Acad Sci USA 96:1732-1737; Nuclear receptor coactivator 1 (Ncoal), Qi C, et al. (1999) Proc Natl Acad Sci USA  
30 96:1585-1590; Decay accelerating factor 1 (Daf1), Sun X, et al. (1999) Proc Natl Acad Sci USA 1999, 96:628-633; Necdin (Ndn), Tsai TF, et al. (1999) Nat Genet 22:15-16; Relaxin (Rln); Zhao L, et al. (1999) Endocrinology 140:445-453; Adenylyl cyclase 8



- (Adcy8), Abdel-Majid RM, et al. (1998) *Nat Genet* 19:289-291; Leukemia inhibitory factor (Lif), Bugga L, et al. (1998) *J Neurobiol* 36:509-524; Lectin, galactose binding, soluble 3 (Lgals3) and Lgals1, Calnot C, et al. (1998) *Dev Dyn* 211:306-313; Urokinase plasminogen activator receptor (Plaur) Carmeliet P, et al. (1998) *J Cell Biol* 140:233-245; Nitric oxide synthase 1, neuronal (Nos1), Chao DS, et al. (1998) *J Neurochem* 71:784-789; Homeo box A7 (Hoxa7), Chen F, et al. (1998) *Mech Dev* 77:49-57; Myosin light chain, phosphorylatable, cardiac ventricles (MyIpc) Chen J, et al. (1998) *J Biol Chem* 273:1252-1256; Homeo box B7 (Hoxb7), Chen F, et al. (1998) *Mech Dev* 77:49-57; Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha and beta (Nfkb1a and Nfkb1b), Cheng JD, et al. (1998) *J Exp Med* 186:1055-1062; Enolase 1, alpha non-neuron (Eno1), Couldrey C, et al. (1998) *Dev Dyn* 212:284-292; Xeroderma pigmentosum, complementation group A (Xpa), De Vries A, et al. (1998) *Exp Eye Res* 1998, 67:53-59; Von Willebrand factor homolog (Vwf), Denis C, et al. (1998) *Proc Natl Acad Sci USA* 95:9524-9529; Lysosomal acid lipase 1 (Lip1), Du H, et al. (1998) *Hum Mol Genet* 7:1347-1354; UNC-5 homolog (*C. elegans*) 3 (Unc5h3), Eisenman LM, et al. (1998) *J Comp Neurol* 394:106-117; Protein phosphatase 1, regulatory (inhibitor) subunit 1B (Ppp1r1b), Fienberg AA, et al. (1998) *Science* 281:838-842; Myelin-associated glycoprotein (Mag) Fujita N, et al. (1998) *J Neurosci* 18:1970-1978; Paraoxonase 1 (Pon1), Furlong CE, et al. (1998) *Neurotoxicology* 19:645-650; Brain derived neurotrophic factor (Bdnf), Garek RR, et al. (1998) *Laryngoscope* 108:671-678; Neurotrophin 3 (Ntf3), Garek RR, et al. (1998) *Laryngoscope* 108:671-678; Myoglobin (Mb); Garry DJ, et al. (1998) *Nature* 395:905-908; Opioid receptor, mu (Oprm), Gaveriaux-Ruff C, et al. (1998) *Proc Natl Acad Sci USA* 95:6326-6330; Neuropeptide Y (Npy), Hollopeter G, et al. (1998) *Int J Obes Relat Metab Disord* 22:506-512; Procollagen, type I, alpha 1 (Col1a1) Hormuzdi SG, et al. (1998) *Mol Cell Biol* 18:3368-3375; Centromere autoantigen B (Cenpb), Hudson DF, et al. (1998) *J Cell Biol* 141:309-319; Oculocerebrorenal syndrome of Lowe (ocrl), Janne PA, et al. (1998) *J Clin Invest* 101:2042-2053; arachidonate 12-lipoxygenase (Alox12) Johnson EN, et al. (1998) *Proc Natl Acad Sci USA* 95:3100-3105; H19 fetal liver mRNA (H19), Jones BK, et al. (1998) *Genes Dev* 12:2200-2207; Hepatocyte nuclear factor 3 gamma (winged helix transcription factor) (Hnf3g), Kaestner KH, et al. (1998) *Mol Cell Biol* 18:4245-4251; Bone morphogenetic protein 2 (Bmp2) / Bone

- morphogenetic protein 7 (Bmp7) and Bmp5, Katagiri T, et al. (1998) *Dev Genet* 22:340–348; Intercellular adhesion molecule (Icam1), Ley K, et al. (1998) *Circ Res* 83:287–294; Glutamyl aminopeptidase (Enpep), Lin Q, et al. *J Immunol* 1998, 160:4681–4687; Prion protein (Prnp), Lipp HP, et al. *Behav Brain Res* 1998, 95:47–54; RAB3A, member RAS oncogene family (Rab3a), Lonart G, et al. *Neuron* 1998, 21:1141–1150; Potassium voltage gated channel, shaker related subfamily, member 4 (Kcna4), London B, et al., *J Physiol (Lond)* 1998, 509:171–182; Apurinic/aprimidinic endonuclease (Apex), Ludwig DL, et al. *Mutat Res* 1998, 409:17–29; T-cell receptor gamma, variable 5 (Tcrg-V5), Mallick-Wood CA, et al. *Science* 1998, 279:1729–1733; Nuclear, factor, erythroid derived 2, like 2 (Nfe2l2), Martin F, et al. *Blood* 1998, 91:3459–3466; Interleukin 13 (Il13), McKenzie GJ, et al. *Curr Biol* 1998, 8:339–340; Sorbitol dehydrogenase 1 (Sdh1), Ng T, et al. *Diabetes* 1998, 47:961–966; Guanine nucleotide binding protein, alpha 11 (Gna11), Offermanns S, et al. *EMBO J* 1998, 17:4304–4312; Estrogen receptor alpha (Estra), Ogawa S, et al. *Endocrinology* 1998, 139:5070–5081; Integrin beta 2 (Itgb2), Intracellular adhesion molecule (Icam1) and CD34 antigen, Oliveira-dos-Santos AJ, et al. *Eur J Immunol* 1998, 28:2882–2892; Angiotensin receptor 1b (Agtr1b), Oliverio MI, et al. *Proc Natl Acad Sci USA* 1998, 95:15496–15501; Complement factor B (factor B), Pekna M, et al. *Scand J Immunol* 1998, 47:375–380; Centromere autoantigen B (Cenpb), Perez-Castro AV, et al. *Dev Biol* 1998, 201:135–143; Procollagen, type V, alpha 2 (Col5a2) / Fibrillin 1 (Fbn1), Phelps RG, et al. *Mol Med* 1998, 4:356–360; Plasminogen activator inhibitor, type 1 (Planh1), Pinsky DJ, et al. *J Clin Invest* 1998, 102:919–928, Carmeliet P, et al. *J Clin Invest* 1993, 92:2756–2760; Placentae and embryos oncofetal gene (Pem), Pitman JL, et al. *Dev Biol* 1998, 202:196–214; Postmeiotic segregation increased 1 (*S. cerevisiae*) (Pms1), Prolla TA, et al. *Nat Genet* 1998, 18:276–278; Prion protein, structural locus (Prn-p), Prusiner SB, et al. *Proc Natl Acad Sci USA* 1998, 95:10608–10612, Lledo P-M, et al. *Proc Natl Acad Sci U S A* 1996, 93:2403–2407, Sailer A, et al. *Cell* 1994, 77:967–968, Weissmann C, et al. *Philos Trans R Soc Lond [Biol]* 1994, 343:431–433, Bueler H, et al. *Cell* 1993, 73:1337–1347, Weissmann C, et al. *Intervirology* 1993, 35:164–175; NAD (P)H:quinone oxidoreductase, Radjendirane V, et al. *J Biol Chem* 1998, 273:7382–7389; Alpha tropomyosin (Tpm1), Rethinasamy P, et al. *Circ Res* 1998, 82:116–123; Goosecoid and Goosecoid-like (Gsc1), Wakamiya M, et al. *Hum Mol Genet* 1998, 7:1835–1840, Saint-Pore B, et al. *Hum Mol Genet* 1998,

- 7:1841-1849; Schlafen 1 (Slfn1); Schwarz DA, et al. *Immunity* 1998, 9:657-668; Nuclear factor, erythroid derived 2, ubiquitous (Mafk), Shavit JA, et al. *Genes Dev* 1998, 12:2164-2174; Microphthalmia-associated transcription factor (Mitf), Smith SB, et al. *Exp Eye Res* 1998, 66:403-410; Bone morphogenetic protein 6 (Bmp6), Solloway MJ, et al. *Dev Genet* 1998, 22:321-339; Phosphatidylinositol glycan, class A (Piga), Takahama Y, et al. *Eur J Immunol* 1998, 28:2159-2166; Paired-related homeobox 2 (Prx2), Berge D, et al. *Development* 1998, 125:3831-3842; Prostaglandin E receptor EP1 subtype (Ptgerp1), Ushikubi F, et al. *Nature* 1998, 395:281-284; Immunoglobulin kappa chain complex (Igk), van der Stoep N, et al. *Immunity* 1998, 8:743-750; Adenine phosphoribosyl transferase (Aprt), Van Sloun PP, et al. *Nucleic Acids Res* 1998, 26:4888-4894; Microtubule associated protein 4 (Mtap4), Voss AK, et al. *Dev Dyn* 1998, 212:258-266; ; 3-hydroxy-3-methylglutaryl-coenzyme A lyase (Hmgcl), Wang SP, et al. *Hum Mol Genet* 1998, 7:2057-2062; Fibroblast growth factor receptor 4 (Fgfr4), Weinstein M, et al. *Development* 1998, 125:3615-3623; Hepsin (Hpn), Wu Q, et al. *J Clin Invest* 1998, 101:321-326; Small inducible cytokine A11 (Scyal1), Yang T, et al. *Blood* 1998, 92:3912-3923; Small nuclear ribonucleoprotein N (Snrpn), Yang T, et al. *Nat Genet* 1998, 19:25-31; DNA fragmentation factor, alpha subunit (Dffa), Zhang J, et al. *Proc Natl Acad Sci USA* 1998, 95:12480-12485; Early growth response 1 (Egr1), Zheng D, et al. *Neuroscience* 1998, 83:251-258; Early growth response 1 (Egr1) / Hormone receptor (Hmr), Zheng D, et al. *Neuroscience* 1998, 83:251-258; Hemochromatosis (Hfe), Zhou XY, et al. *Proc Natl Acad Sci USA* 1998, 95:2492-2497; Alpha tropomyosin (Tpm1), Blanchard EM, et al. *Circ Res* 1997, 81:1005-1010; tRNA phosphoserine (Trsp), Bosl MR, et al. *Proc Natl Acad Sci U S A* 1997, 94:5531-5534; Angiotensin receptor 1b (Atr1b), Chen X, et al. *Am J Physiol* 1997, 272:F299-F304; Xeroderma pigmentosum, complementation group C (Xpc), Cheo DL, et al. *Mut Res* 1997, 374:1-9; B cell leukemia/lymphoma 6 (Bcl6), Dent AL, et al. *Science* 1997, 276:589-592; Fumarylacetoacetate hydrolase (Fah) / 4-hydroxyphenylpyruvic acid dioxygenase (Hpd), Endo F, et al. *J Biol Chem* 1997, 272:24426-24432; N-methylpurine-DNA glycosylase (Mpg), Engelward BP, et al. *Proc Natl Acad Sci USA* 1997, 94:13087-13092; Interleukin 1 receptor, type 1 (Il1r1), Glaccum MB, et al. *J Immunol* 1997, 159:3364-3371; N-methylpurine-DNA glycosylase (Mpg), Hang B, et al. *Proc Natl Acad Sci USA* 1997, 94:12869-12874; Gamma-aminobutyric acid (GABA-

- A) receptor, subunit alpha 6 (Gabra6), Homanics GE, et al. *Mol Pharmacol* 1997, 51:588-596; Superoxide dismutase 2, mitochondrial (Sod2), Huang TT, et al. *Arch Biochem Biophys* 1997, 344:424-432; Interleukin 11 receptor, alpha chain 1 (Il11ra1), Nandurjar HH, et al. *Blood* 1997, 90:2148-2159; Alkaline phosphatase 5 (Akp5),
- 5 Narisawa S, et al. *Dev Dyn* 1997, 208:432-446; GATA-binding protein 4 (Gata4), Narita N, et al. *Development* 1997, 124:3755-3764; Lymphocyte protein tyrosine kinase (Lck) / Fyn protooncogene (Fyn); Page ST, et al. *Eur J Immunol* 1997, 27:554-562; P glycoprotein 3 (Pgy3), Schinkel AH, et al. *Proc Natl Acad Sci U S A* 1997, 94:4028-4033; P glycoprotein 1 (Pgy1) / P glycoprotein 3 (Pgy3), Schinkel AH, et al., *Proc Natl*
- 10 *Acad Sci U S A* 1997, 94:4028-4033; Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. *J Neurosci Methods* 1997, 71:29-41; T cell receptor gamma, variable 4 (Tcrg-V4), Sunaga S, et al. *J Immunol* 1997, 158:4223-4228; Ia-associated invariant chain (p31 form) (Ii), Takaesu NT, et al. *J Immunol* 1997, 158:187-199; Solute carrier family 18 (vesicular monoamine), member 2 (Slc18a2), Takahashi N, et al. *Proc*
- 15 *Natl Acad Sci USA* 1997, 94:9938-9943; Matrix metalloproteinase 7 (Mmp7), Wilson CL, et al. *Proc Natl Acad Sci U S A* 1997, 94:1402-1407; Formin (Fmn), Wynshaw-Boris A, et al. *Mol Med* 1997, 3:372-384; Synaptophysin (Syp), Arrandale JM, et al. *J Biol Chem* 1996, 271:21353-21358; Transformation related protein 53 (Trp53), Boehme SA, et al. *J Immunol* 1996, 156:4075-4078; Neuronal nitric oxide synthase (Nos1),
- 20 Burnett AL, et al. *Mol Medicine* 1996, 2:288-296; Eph receptor A2 (Epha2), Chen J, et al. *Oncogene* 1996, 12:979-988; Urokinase plasminogen activator receptor (Plaur); Dewerchin M, et al. *J Clin Invest* 1996, 97:870-878; Growth differentiation factor 9 (Gdf9), Dong J, et al. *Nature* 1996, 383:531-535; Externally regulated phosphatase (Ptpn16), Dorfman K, et al. *Oncogene* 1996, 13:925-931; Tenascin C (Tnc), Forsberg E,
- 25 et al. *Proc Natl Acad Sci U S A* 1996, 93:6594-6599; Integrin alpha 1 (Itga1), Gardner H, et al. *Dev Biol* 1996, 175:301-313; FBJ osteosarcoma oncogene B (Fosb), Gruda MC, et al. *Oncogene* 1996, 12:2177-2185; Breast cancer 1 (Brca1), Hakem R, et al. *Cell* 1996, 85:1009-1023; Megakaryocyte-associated tyrosine kinase (Matk), Hamaguchi I, et al. *Biochem Biophys Res Commun* 1996, 224:172-179; Apolipoprotein B editing
- 30 complex 1 (Apobec1), Hirano K-I, et al. *J Biol Chem* 1996, 271:9887-9890; Carboxyl ester lipase, Howles PN, et al. *J. Biol. Chem.* 1996, 271:7196-7202; Nuclear factor, erythroid derived 2, ubiquitous (Nfe2u), Kotkow KJ, Orkin SH. *Proc Natl Acad Sci U S*

- A 1996, 93:3514–3518; Retinoid X receptor gamma (Rxrg), Krezel W, et al. *Proc Natl Acad Sci U S A* 1996, 93:9010–9014; Early growth response 1 (Egr1), Lee SL, et al. *Mol Cell Biol* 1996, 16:4566–4572; Adrenergic receptor, alpha 2b (Adra2b), Link RE, et al. *Science* 1996, 273:803–805; Angiotensin receptor 1a (Agtr1a), Matsusaka T, et al. *J Clin Invest* 1996, 98:1867–1877; Interleukin 3 receptor, beta chain 1 (Il3rb2), Nicola NA, et al. *Blood* 1996, 87:2665–2674; Transformation related protein 53 (Trp53), Ohashi M, et al. *Jpn J Cancer Res* 1996, 87:696–701; Leukemia-associated gene (Lag), Schubart U K, et al. *J Biol Chem* 1996, 271:14062–14066; Glutathione peroxidase 1 (Gpx1), Spector A, et al. *Exp Eye Res* 1996, 62:521–540; Arachidonate 12-lipoxygenase, leukocyte (Alox12l), Sun D, Funk CD. *J Biol Chem* 1996, 271:24055–24062; Complement component 3 (C3), Sylvestre D, et al. *J Exp Med* 1996, 184:2385–2392; CD30 antigen (Cd30), Texido G, et al. *Eur J Immunol* 1996, 26:1966–1969; Fc receptor, IgE, low affinity II, alpha polypeptide (Fcer2a), Immunoglobulin heavy chain 5 (delta-like heavy chain) (Igh-5), Interleukin 4 (IL4) and Terminal-deoxynucleotidyl transferase, Texido G, et al. *Eur J Immunol* 1996, 26:1966–1969; Apolipoprotein A-II (Apoa2), Weng W, Breslow JL. *Proc Natl Acad Sci U S A* 1996, 93:14788–14794; Amyloid beta (A4) precursor protein (App), Zheng H, et al. *Ann N Y Acad Sci* 1996, 777:421–426; cAMP responsive element binding protein 1 (Creb1), Blendy JA, et al. *Brain Res* 1995, 681:8–14; Bradykinin receptor, beta 2 (Bdkrb2), Borkowski JA, et al. *J Biol Chem* 1995, 270:13706–13710; Growth factor response protein (Gfrp), Crawford PA, et al. *Mol Cell Biol* 1995, 15:4331–4336; Ciliary neurotrophic factor (Cntf), de Chiara TM, et al. *Cell* 1995, 83:313–322; Cyclin dependent kinase inhibitor 1A (P21) (Cdkn1a); Deng C, et al. *Cell* 1995, 82:675–684; Granzyme A (Gzma), Ebnet K, et al. *EMBO J* 1995, 14:4230–4239; Very low density lipoprotein receptor (Vldlr), Frykman PK, et al. *Proc Natl Acad Sci U S A* 1995, 92:8453–8457; Apolipoprotein E (ApoE) / Apolipoprotein A-I (Apoa1), Goodrum JF, et al. *J Neurobiol* 1995, 64:408–416; Nitric oxide synthase 1, neuronal (Nos1), Ichinose F, et al. *Anesthesiology* 1995, 83:101–108; Nitric oxide synthase 2, inducible, macrophage (Nos2), Laubach VE, et al. *Proc Natl Acad Sci U S A* 1995, 92:10688–10692; Peroxisome proliferator activated receptor alpha (Ppara), Lee SS-T, et al. *Mol Cell Biol* 1995, 15:3012–3022; Growth factor response protein (Gfrp), Lee SL, et al. *Science* 1995, 269:532–535; H19 fetal liver mRNA (H19) / Insulin-like growth factor 2 (Igf2), Leighton PA, et al. *Nature* 1995, 375:34–39; Retinoic

- acid receptor beta (RAR-beta), Luo J, et al. *Mech Dev* 1995, 53:61-71; Metallothionein 1 (Mt1) / Metallothionein 2 (Mt2), Philcox JC, et al. *Biochem J* 1995, 308:543-546; Heme oxygenase (decycling) 2 (Hmox2), Poss KD, et al. *Neuron* 1995, 15:867-873; H1-0 histone (H1fv), Sirotkin AM, et al. *Proc Natl Acad Sci U S A* 1995, 92:6434-6438;
- 5 Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. *Biochim Biophys Acta* 1995, 1230:130-138; Tenascin C (Tnc), Steindler DA, et al. *J Neurosci* 1995, 15:1971-1983; Ia-associated invariant chain (Ii), Takaesu NT, et al. *Immunity* 1995, 3:385-396; Neuroblastoma ras oncogene (Nras), Umanoff H, et al. *Proc Natl Acad Sci USA* 1995, 92:1709-1713; Receptor-associated protein of the synapse, 43 kDa (Rapsn),
- 10 Willnow TE, et al. *Proc Natl Acad Sci USA* 1995, 92:4537-4541; Vitronectin (Vtn), Zheng X, et al. *Proc Natl Acad Sci U S A* 1995, 92:12426-12430; Preproacrosin (Acr), Baba T, et al. *J Biol Chem* 1994, 269:31845-31849; Vimentin (Vim), Colucci GE, et al. *Cell* 1994, 79:679-694; Tumor necrosis factor receptor superfamily, member 1b (Tnfrsf1b), Erickson SL, et al. *Nature* 1994, 372:560-563; Cellular retinoic acid binding
- 15 protein I (Crabp1), Gorry P, et al. *Proc Natl Acad Sci USA* 1994, 91:9032-9036; cAMP responsive element binding protein 1 (Creb1), Hummler F, et al. *Proc Natl Acad Sci USA* 1994, 91:5647-5651; Pore forming protein (Pfp), Kagi D, et al. *Nature* 1994, 369:31-37; Src-related kinase lacking C-terminal regulatory tyrosine and N-terminal myristylation sit (Srms), Kohmura N, et al. *Mol Cell Biol* 1994, 14:6915-6925; CD3
- 20 polypeptide zeta (Cd3z) / Solute carrier family 22, member 1 (Slc22a1), Koyasu S, et al. *EMBO J* 1994, 13:784-797; Pore forming protein (Pfp), Lowin B, et al. *Proc Natl Acad Sci USA* 1994, 91:11571-11575; Retinoic acid receptor, beta (Rarb), Retinoic acid receptor beta2 (RARbeta2), Mendelsohn C, et al. *Dev Biol* 1994, 166:246-258; Transthyretin (Tr), Palha JA, et al. *J Biol Chem* 1994, 269:33135-33139; Procollagen,
- 25 type X, alpha 1 (Col10a1), Rosati R, et al. *Nature Genet* 1994, 8:129-135; P glycoprotein 3 (Pgy3), Schinkel AH, et al. *Cell* 1994, 77:491-502; Yamaguchi sarcoma viral (v-yes) oncogene homolog (Yes), Stein PL, et al. *Genes Dev* 1994, 8:1999-2007; Fc receptor, IgE, high affinity II, alpha polypeptide (Fcer2a), Stief A, et al. *J Immunol* 1994, 152:3378-3390; Pore forming protein (Pfp), Walsh CM, et al. *Proc Natl Acad Sci*
- 30 *USA* 1994, 91:10854-10858; CD2 antigen (Cd2), Evans CF, et al. *J Immunol* 1993, 151:6259-6264; Mannose-6-phosphate receptor, cation dependent (M6pr), Koster A, et al. *EMBO J* 1993, 12:5219-5223; Retinoic acid receptor, alpha (Rara), Li E, et al. *Proc*

- Natl Acad Sci USA 1993, 90:1590-1594, Lufkin T, et al. Proc Natl Acad Sci USA 1993, 90:7225-7229; Retinoic acid receptor, gamma (Rarg), Lohnes D, et al. Cell 1993, 73:643-658; Tumor necrosis factor receptor 1 (TNF-R-1) (Tnfr1), Pfeffer K, et al. Cell 1993, 73:457-467; Lectin, galactose binding, soluble 1 (Lgals1), Poirier F, Robertson EJ.
- 5 Development 1993, 119:1229-1236; Synapsin I (Syn1), Rosahl TW, et al. Cell 1993, 75:661-670; Tumor necrosis factor receptor 1 (Tnfr1), Rothe J, et al. Nature 1993, 364:798-802; Beta-2 microglobulin (B2m), Correa I, et al. Proc Natl Acad Sci USA 1992, 89:653-657; CD2 antigen (Cd2), Killeen N, et al. EMBO J 1992, 11:4329-4336; Apolipoprotein E (ApoE), Piedrahita JA, et al. Proc Natl Acad Sci USA 1992, 89:4471-
- 10 4475; Myogenic differentiation 1 (Myod1), Rudnicki MA, et al. Cell 1992, 71:383-390; Tenascin C (Tnc), Saga Y, et al. Genes Dev 1992, 6:1821-1831; Beta-2 microglobulin (B2m), Sanjuan N, et al. J Virol 1992, 66:4587-4590; Neuroblastoma myc-related oncogene 1 (Nmyc1), Stanton BR, et al. Genes Dev 1992, 6:2235-2247; and Hemoglobin alpha chain complex (Hba), Popp RA, et al. Genetics 1983, 105:157-167.
- 15 Some preferred single-copy, non-essential genes with no phenotypes of the present invention include, but are not limited to, the following: Moesin (Msn), Doi Y., et al., J Biol Chem 1999, 274:2315-2321; Plasminogen activator inhibitor, type II (Planh2) and Planh1, Dougherty K.M., Proc Natl Acad Sci USA 1999, 96:686-691; Nuclear receptor coactivator 1 (Nco1), Qi C, et al. (1999) Proc Natl Acad Sci USA 96:1585-
- 20 1590; Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha and beta (Nfkb1a and Nfkb1b), Cheng JD, et al. (1998) J Exp Med 6:1055-1062; H19 fetal liver mRNA (H19), Jones BK, et al. (1998) Genes Dev 12:2200-2207; Prion protein (Prnp), Lipp HP, et al. Behav Brain Res 1998, 95:47-54; Centromere autoantigen B (Cenpb), Perez-Castro AV, et al. Dev Biol 1998, 201:135-143; Placentae and embryos oncofetal gene (Pem), Pitman JL, et al. Dev Biol 1998, 202:196-214; Externally
- 25 regulated phosphatase (Ptpn16), Dorfman K, et al. Oncogene 1996, 13:925-931; Transformation related protein 53 (Trp53), Ohashi M, et al. Jpn J Cancer Res 1996, 87:696-701; H1-0 histone (H1fv), Sirotkin AM, et al. Proc Natl Acad Sci U S A 1995, 92:6434-6438; Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al.
- 30 Biochim Biophys Acta 1995, 1230:130-138; Neuroblastoma ras oncogene (Nras), Umanoff H, et al. Proc Natl Acad Sci USA 1995, 92:1709-1713; Vitronectin (Vtn), Zheng X, et al. Proc Natl Acad Sci U S A 1995, 92:12426-12430; Vimentin (Vim),

Colucci GE, et al. Cell 1994, 79:679-694; Cellular retinoic acid binding protein I (Crabp1), Gorry P, et al. Proc Natl Acad Sci USA 1994, 91:9032-9036; Retinoic acid receptor beta2 (RARbeta2), Mendelsohn C, et al. Dev Biol 1994, 166:246-258; Retinoic acid receptor, alpha (Rara), Li E, et al. Proc Natl Acad Sci USA 1993, 90:1590-1594,  
 5 Lufkin T, et al. Proc Natl Acad Sci USA 1993, 90:7225-7229; Lectin, galactose binding, soluble 1 (Lgals1), Poirier F, Robertson EJ. Development 1993, 119:1229-1236; Myogenic differentiation 1 (Myod1), Rudnicki MA, et al. Cell 1992, 71:383-390; and Tenascin C (Tnc), Saga Y, et al. Genes Dev 1992, 6:1821-1831.

In view of the guidance of the present specification, one of ordinary skill in the  
 10 art can select similar, suitable, single-copy, non-essential genes in mice and other cell types/organisms.

### Assembly of Targeting Cassettes

The targeting cassettes described herein can be constructed utilizing  
 15 methodologies known in the art of molecular biology (see, for example, Ausubel or Maniatis) in view of the teachings of the specification. As described above, the targeting constructs are assembled by inserting, into a suitable vector backbone, polynucleotides encoding a reporter, such as a light-generating protein, *e.g.*, a luciferase gene, operably linked to a promoter of interest; a sequence encoding a positive selection marker; and,  
 20 optionally a sequence encoding a negative selection marker. In addition, the targeting cassette contains insertion sites such that sequences targeting a single-copy, non-essential gene can be readily inserted to flank the sequence encoding positive selection marker and luciferase-encoding sequence.

A preferred method of obtaining polynucleotides, suitable regulatory sequences  
 25 (*e.g.*, promoters) is PCR. General procedures for PCR as taught in MacPherson et al., PCR: A PRACTICAL APPROACH, (IRL Press at Oxford University Press, (1991)). PCR conditions for each application reaction may be empirically determined. A number of parameters influence the success of a reaction. Among these parameters are annealing temperature and time, extension time, Mg<sup>2+</sup> and ATP concentration, pH, and the relative  
 30 concentration of primers, templates and deoxyribonucleotides. Exemplary primers are described below in the Examples. After amplification, the resulting fragments can be



detected by agarose gel electrophoresis followed by visualization with ethidium bromide staining and ultraviolet illumination.

In one embodiment, PCR can be used to amplify fragments from genomic libraries. Many genomic libraries are commercially available. Alternatively, libraries  
5 can be produced by any method known in the art. Preferably, the organism(s) from which the DNA is has no discernible disease or phenotypic effects. This isolated DNA may be obtained from any cell source or body fluid (*e.g.*, ES cells, liver, kidney, blood cells, buccal cells, cerviovaginal cells, epithelial cells from urine, fetal cells, or any cells present in tissue obtained by biopsy, urine, blood, cerebrospinal fluid (CSF), and tissue  
10 exudates at the site of infection or inflammation). DNA is extracted from the cells or body fluid using known methods of cell lysis and DNA purification. The purified DNA is then introduced into a suitable expression system, for example a lambda phage.

Another method for obtaining polynucleotides, for example, short, random nucleotide sequences, is by enzymatic digestion. As described below in the Examples,  
15 short DNA sequences generated by digestion of DNA from vectors carrying genes encoding luciferase (yellow green or red).

Polynucleotides are inserted into vector genomes using methods known in the art. For example, insert and vector DNA can be contacted, under suitable conditions, with a restriction enzyme to create complementary or blunt ends on each molecule that can pair  
20 with each other and be joined with a ligase. Alternatively, synthetic nucleic acid linkers can be ligated to the termini of a polynucleotide. These synthetic linkers can contain nucleic acid sequences that correspond to a particular restriction site in the vector DNA. Other means are known and, in view of the teachings herein, can be used.

25 The final constructs can be used immediately (*e.g.*, for introduction into ES cells), or stored frozen (*e.g.*, at -20°C) until use. Preferably, the constructs are linearized prior to use, for example by digestion with suitable restriction endonucleases.

### Transgenic Animals

The targeting constructs containing the light generating protein coding sequences (e.g., luciferase genes) are introduced into a pluripotent cell (e.g., ES cell, Robertson, E. J., In: Current Communications in Molecular Biology, Capecchi, M. R. (ed.), Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1989), pp. 39-44). Suitable ES cells may be derived or isolated from any species or from any strain of a particular species. Although not required, the pluripotent cells are typically derived from the same species as the intended recipient. ES cells may be obtained from commercial sources, from International Depositories (e.g., the ATCC) or, alternatively, may be obtained as described in Robertson, E. J., *supra*. Examples of clonally-derived ES cell lines include 129/SVJ ES cells, RW-4 and C57BL/6 ES cells (Genome Systems, Inc.).

ES cells are cultured under suitable conditions, for example, as described in Ausubel et al., section 9.16, *supra*. Preferably, ES cells are cultured on stromal cells (such as STO cells (especially SNC4 STO cells) and/or primary embryonic fibroblast cells) as described by E. J. Robertson, *supra*, pp 71-112. Culture media preferably includes leukocyte inhibitory factor ("lif") (Gough, N. M. et al., Reprod. Fertil. Dev. 1:281-288 (1989); Yamamori, Y. et al., Science 246:1412-1416 (1989), which appears to help keep the ES cells from differentiating in culture. Stromal cells transformed with the gene encoding lif can also be used.

The targeting constructs are introduced into the ES cells by any method which will permit the introduced molecule to undergo recombination at its regions of homology, for example, micro-injection, calcium phosphate transformation, or electroporation (Toneguzzo, F. et al., Nucleic Acids Res. 16:5515-5532 (1988); Quillet, A. et al., J. Immunol. 141:17-20 (1988); Machy, P. et al., Proc. Natl. Acad. Sci. (U.S.A.) 85:8027-8031 (1988)). The construct to be inserted into the ES cell must first be in the linear form. Thus, if the knockout construct has been inserted into a vector as described above, linearization is accomplished by digesting the DNA with a suitable restriction endonuclease selected to cut only within the vector sequence and not within the knockout construct sequence. If the ES cells are to be electroporated to insert the construct, the ES cells and construct DNA are exposed to an electric pulse using an electroporation machine and following the manufacturer's guidelines for use. After electroporation, the ES cells are typically allowed to recover under suitable incubation conditions. The cells

are then cultured under conventional conditions, as are known in the art, and screened for the presence of the construct.

Screening and selection of those cells into which the targeting construct has been integrated can be achieved using the positive selection marker and/or the negative  
5 selection marker in the construct. In preferred embodiments, the construct contains both positive and negative selection markers. In one aspect, methods which rely on expression of the selection marker are used, for example, by adding the appropriate substrate to select only those cells which express the product of the positive selection marker or to eliminate those cells expressing the negative selection marker. For  
10 example, where the positive selection marker encodes neomycin resistance, G418 is added to the transformed ES cell culture media at increasing dosages. Similarly, where the negative selection marker is used, a suitable substrate (*e.g.*, gancyclovir if the negative selection marker encodes HSV-TK) is added to the cell culture. Either before or after selection using the appropriate substrate, the presence of the positive and/or  
15 negative selection markers in a recipient cell can also be determined by others methods, for example, hybridization, detection of radiolabelled nucleotides, PCR and the like. In preferred embodiments, cells having integrated targeting constructs are first selected by adding the appropriate substrate for the positive and/or negative selection markers. Cells that survive the selection process are then screened by other methods, such as PCR or  
20 Southern blotting, for the presence of integrated sequences.

After suitable ES cells containing the construct in the proper location have been identified, the cells can be inserted into an embryo, preferably a blastocyst. The blastocysts are obtained by perfusing the uterus of pregnant females. In one embodiment, the blastocysts are obtained from, for example, the FVB/N strain of mice and the ES cells  
25 are obtained from, for example, the C57BL/6 strain of mice. Suitable methods for accomplishing this are known to the skilled artisan, and are set forth by, *e.g.*, Bradley et al., (1992) *Biotechnology*, 10:534-539. Insertion into the embryo may be accomplished in a variety of ways known to the skilled artisan, however a preferred method is by microinjection. For microinjection, about 10-30 ES cells are collected into a micropipet  
30 and injected into embryos that are at the proper stage of development to permit integration of the foreign ES cell containing the construct into the developing embryo.

The suitable stage of development for the embryo used for insertion of ES cells is species dependent, in mice it is about 3.5 days.

While any embryo of the right stage of development is suitable for use, it is preferred that blastocysts are used. In addition, preferred blastocysts are male and, 5 furthermore, preferably have genes encoding a coat color that is different from that encoded by the genes ES cells. In this way, the offspring can be screened easily for the presence of the knockout construct by looking for mosaic coat color (indicating that the ES cell was incorporated into the developing embryo). Thus, for example, if the ES cell line carries the genes for black fur, the blastocyst selected will carry genes for white or 10 brown fur.

After the ES cell has been introduced into the blastocyst, the blastocyst is typically implanted into the uterus of a pseudopregnant foster mother for gestation. . . . Pseudopregnant females are prepared by mating with vasectomized males of the same species and successful implantation usually must occur within about 2-3 days of mating.

15 Offspring are screened initially for mosaic coat color where the coat color selection strategy has been employed. Southern blots and/or PCR may also be used to determine the presence of the sequences of interest. Mosaic (chimeric) offspring are then bred to each other to generate homozygous animals. Homozygotes and heterozygotes may be identified by Southern blotting of equivalent amounts of genomic DNA from 20 mice that are the product of this cross, as well as mice that are known heterozygotes and wild type mice. Alternatively, Northern blots can be used to probe the mRNA to identify the presence or absence of transcripts encoding either the replaced gene, the light generating protein coding sequence (e.g., luciferase gene), or both. In addition, Western blots can be used to assess the level of expression of the luciferase protein with an 25 antibody against the luciferase gene product. Finally, in situ analysis (such as fixing the cells and labeling with antibody) and/or FACS (fluorescence activated cell sorting) analysis of various cells from the offspring can be conducted using suitable (e.g., anti-luciferase) antibodies to look for the presence or absence of the targeting construct.

In one embodiment of the present invention, the animals are from the C57BL/6 30 mouse strain. This strain develops a variety of tumors and has been used to develop a number of tumor cells lines, for example, B16 melanoma cells (including, B16F10, B16D5, and B16F1), Lewis lung carcinoma cells (including, LLC, LLC-h59), T241

mouse fibrosarcoma cells, RM-1 and pTC2 mouse prostate cancer cells, and MCA207 mouse sarcoma cells. These cell lines have been extensively used for *in vivo* tumor biology studies after injection into C57BL/6 mice. The generated targeted transgenic mice in the Examples are in C57BL/6 genetic background and these animals are suitable for injection or implantation of such tumor cells, as well as other tumor cells described in literature that are immunocompatent for C57BL/6 mice. Thus, the transgenic animals can then be used, for example, to monitor, *in vivo*, tumor progression (*e.g.*, growth) and the efficacy of therapies on tumor regression. For example, where the transgenic animal is tumor-susceptible, it is monitored for expression of a reporter, *e.g.*, luciferase, which is indicative of tumorigenesis and/or angiogenesis. The monitoring of expression of luciferase reporter expression cassettes using non-invasive whole animal imaging has been described (Contag, C. et al, U.S. Patent No. 5,650,135, July 22, 1997; Contag, P., et al, *Nature Medicine* 4(2):245-247, 1998; Contag, C., et al, *OSA TOPS on Biomedical Optical Spectroscopy and Diagnostics* 3:220-224, 1996; Contag, C.H., et al, *Photochemistry and Photobiology* 66(4):523-531, 1997; Contag, C.H., et al, *Molecular Microbiology* 18(4):593-603, 1995). Such imaging typically uses at least one photo detector device element, for example, a charge-coupled device (CCD) camera.

The transgenic animals described herein can also be used to determine the effect of an analyte (*e.g.*, therapy), for example on tumor progression where the promoter induces light generating protein (*e.g.*, luciferase) expression when a tumor develops. Methods of administration of the analyte include, but are not limited to, injection (subcutaneously, epidermally, intradermally), intramucosal (such as nasal, rectal and vaginal), intraperitoneal, intravenous, oral or intramuscular. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. For example, the analyte of interest can be administered over a range of concentration to determine a dose/response curve. The analyte may be administered to a series of test animals or to a single test animal (given that response to the analyte can be cleared from the transgenic animal).

The following examples are intended only to illustrate the present invention and should in no way be construed as limiting the subject invention.

## EXAMPLES

### **Example 1 Generating the Targeting Cassette and Vector**

5

#### **A. Creation of the Backbone Vector**

pTK53: The 0.5 kb mouse phosphoglycerate kinase 1 promoter was amplified with PGK primers (PGKF, SEQ ID NO:1:

ATCGAATTCTACCGGGTAGGGGAGGCGCTTT; PGKR, SEQ ID NO:2:

10 GGCTGCAGGTCGAAAGGCCCGGAGATGAGG) using mouse genomic DNA (Genome Systems, Inc., St. Louis, MO) as template. This fragment was then double digested with EcoRI and PstI and cloned into the pKS vector (Stratagene, La Jolla, California) which was linearized with the same enzymes. The neomycin gene was amplified with NeoF (SEQ ID NO:3:

15 ACCTGCAGCCAATATGGGATCGGCCATTGAAC) and NeoR (SEQ ID NO:4: GGATCCGCGGCCGCCCCAGCTGGTTCTTTCCGCCTC) primers using pNTKV1907 (Stratagene) as a template. The 1.1 kb PCR fragment was double digested with PstI and BamHI and cloned into the pKS-PGK vector which was linearized with the same enzymes. This pKS-PGK-Neo vector was used to clone thymidine kinase gene as follows. Primers TKF (SEQ ID NO: 5:

GGATCCTCTAGAGTCGAGCAGTGTGGTTTT) and TKR (SEQ ID NO:6:

GAGCTCCCGTAGTCAGGTTTAGTTCGTCCG) were used to amplify the TK gene from pNTKV1907 (Stratagene). The amplified 2kb fragment was then digested with

25 BamHI and SacI and cloned into pKS-PGK-Neo vector that was linearized with the same enzymes. This constructed vector was designated as pTK. A synthetic linker F5R5 was made after annealing of two primers (forward primer, F5R51, SEQ ID NO: 7:

GTACATTTAAATCCTGCAGG; reverse primer, F5R52, SEQ ID NO:8:

AGCTCCTGCAGGATTTAAAT). This linker was inserted between Asp718I and

HindIII sites of pTK and the new construct was designated pTK5. A second synthetic

30 linker F3R3 was made by annealing of two primers (forward primer, F3R31, SEQ ID NO:9:

GGCCCGGGCTTAATTAATGCATCATATGGTACCGTTTAAACGCGGCCGCAAG

CTTGTCGACGGCGCGCCGGCCGCC; reverse primer, F3R32, SEQ ID NO:10:

GATCGGCCGGCCGGCGCGCCGTCGACAAGCTTGCGGCCGCGTTTAAACGGTACCATATGATGCATTAATTAAGCCCG). ). This linker was inserted between NotI and BamHI sites of pTK and the new construct was designated pTK53. Schematics of the vectors are shown in Figure 1.

5

#### **B. Introduction of Luciferase**

**pTK-LucYG and pTK-LucR:** The yellow green luciferase gene was isolated from pGL3 vector (Promega) as a HindIII-SalI fragment and was cloned into pGK53 that was linearized with the same enzymes. The new construct was designated pTK-LucYG (8931 bp), shown in Figure 2.

10

The red luciferase gene was isolated from pGL3-red vector (Dr. Christopher Contag, Stanford University, Stanford, Calif.) as a HindIII-SalI fragment and was cloned into pGK53 that was linearized with the same enzymes. The new construct was designated pTK-LucR (8931 bp), shown in Figure 2.

15

### **Example 2**

#### **Insertion of Targeting Sequences**

**A. Generation of vitronectin targeting vector:** The targeting construct pTKLR-Vn was generated by inserting vitronectin (VN) DNA sequences into pTK-LucR vector.

20

Vitronectin (VN) is an abundant glycoprotein present in plasma and the extracellular matrix of most tissues. In a previous study, it was shown that heterozygous mice carrying one normal and one null VN allele and homozygous null mice completely deficient in vitronectin demonstrate normal development, fertility, and survival. This suggests that VN is not essential for cell adhesion and migration during normal mouse development (Zheng, X., et al., Proc Natl Acad Sci U S A 1995 92:12426-30). Mouse vitronectin genomic DNA sequence of 5004 bp was obtained from GenBank database (Accession number X72091). Based on this sequence, a 1.63 kb 3'end vitronectin fragment was amplified (reverse primer, VN1R, SEQ ID NO:11: CTGTATTAAATCTGCCACCCCTATTCAGGACAGTAGTC; forward primer, VN1F, SEQ ID NO:12: CCAATGCATCAACCCAGCCAGGAGGAGTGCG) using mouse C57BL/6 genomic DNA as template (Genome Systems, Inc., St. Louis, MO).

25

30

This fragment was digested with *SwaI* and *NsiI* and cloned into pTK-LucR (linearized with *SwaI* and *SbfI*). This construct was designated as pTK-LucR3. Subsequently, a 2.35 kb 5'end vitronectin fragment was amplified (reverse primer, VN2R, SEQ ID NO:13: AACGCGTCGACTTCGGAGATGTTTCGGGGATAACCAGG; forward primer,

5 VN2F, SEQ ID NO:14:

TTGGCGCGCCCCATAGAGAAGAGACACCAAAGGCACGCTC) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *Sall* and *AscI* and cloned into pTK-LucR vector that was linearized with *Sall* and *AscI*. This construct was designated as pTKLR-Vn. Figure 3 shows the restriction map of pTKLR-Vn vector. The  
10 polylinker between the neomycin gene and red luciferase gene is used to insert the VEGF promoter or other promoters of interests. The predicted homologous recombination between pTKLR-Vn and vitronectin gene is illustrated in Figure 3A. Upon insertion of the VEGF-LucR transgene cassette, the endogenous vitronectin gene is destroyed. Figure 3B shows the genomic DNA sequence of VN.

15

**B. Generation of Fos targeting vector:** The targeting construct pTKLG-Fos was generated by inserting FosB DNA sequences into pTK-LucYG vector.

FosB is one of the members of the Fos family. It plays a functional role in transcriptional regulation. It has been shown that FosB mice are born at a normal  
20 frequency, are fertile and present no obvious phenotypic or histologic abnormalities (Gruda et al (1996) *Oncogene* 12:2177-2185). A 28.8 kb genomic region that contains mouse FosB DNA sequence was obtained from GenBank database (Accession number AF093624).

Using this sequence, a 1.71 kb 5'end FosB fragment was amplified (forward  
25 primer, FosB1F, SEQ ID NO:15: CTGTATTAAATCCCGTTTCTCACTGTGCCTGTGTC; reverse primer, FosB1R, SEQ ID NO:16: GTCTCCTGCAGGCTTCCTCCTCCTTGTTTCCTTGCG) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *SwaI* and *SbfI* and cloned into pTK-LucYG vector that was linearized with *SwaI* and *SbfI*. This construct  
30 was designated as pTK-LucYG3. Subsequently, a 1.58 kb 3'end FosB fragment was amplified (forward primer, FosB2F, SEQ ID NO:17: AACGCGTCGACGGATGGGATTGACCCCCAGCCCTC; reverse primer, FosB2R,



SEQ ID NO:18: TTGGCGCGCCCCTTGCCTCCACCTCTCAAATGC) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *Sall* and *Ascl* and cloned into pTK-LucYG vector that was linearized with *Sall* and *Ascl*. This construct was designated as pTKLG-Fos (Figure 4). The polylinker between the neomycin gene and red luciferase gene is used to insert the VEGFR2 promoter (Example 3, Figure 5A), Tie2 promoter (Example 3, Figure 5B), as well as, other promoters of interests. The predicted homologous recombination between the targeting vector bearing the VEGFR2 promoter (Figure 5A) or the Tie2 promoter (Figure 5B) and FosB gene is also illustrated. As shown in the Figures, the VEGFR2-LucYG transgene cassette and Tie2-LucYG transgene cassette is inserted downstream of FosB gene translational stop signal. Therefore, the targeted transgenic mice should still have a functional FosB gene while expressing the transgenes. Figure 4B shows the DNA sequence of FosB.

### Example 3

#### Insertion of Promoter Sequences of Interest

**A. pTKLR-Vn/VEGF:** Mouse VEGF genomic DNA sequence of 2240 bp that contains a partial VEGF promoter region was obtained from GenBank (accession number: U41383). Accordingly, primers were designed to amplify a 0.69 kb (VF1-VR1A; Table 1) and a 0.98 kb fragment (VF2-VR2; Table 1). It was confirmed that each pair of primers can amplify the predicted product using mouse129SvJ genomic DNA as template.

Table 1

Name	Seq Id No	Sequence
VF1	19	ACCTC ACTCT CCTGT CTCCC CTGAT TCCCA A
VR1A	20	GCTCT GGC GG TCACC CCCAA AAGCA
VF2	21	CCCTT TCCAA GACCC GTGCC ATTTG AGC
VR2	22	ACTTT GCCCC TGTCC CTCTC TCTGT TCGC
KF1	23	GCTGC GTCCA GATTT GCTCT CAGAT GCG
KR1	24	TTCTC AGGCA CAGAC TCCTT CTCCG TCCCT
KF2	25	CAGAT GGACG AGAAA ACAGT AGAGG CGTTG GC
KR2	26	GAGGA CTCAG GGCAG AAAGA GAGCG
TF3	27	AGCTT AGCCT GCAAG GGTGG TCCTC ATCG
TF2	28	CAAAT GCACC CCAGA GAACA GCTTA GCCTG C
TR1	29	GCTTT CAACA ACTCA CAACT TTGCG ACTTC CCG

Conditions for PCR amplification are shown in Figure 6. These primers were used for PCR screening of mouse 129/SvJ genomic DNA BAC (bacterial artificial chromosome) library (Genome Systems, Inc., St. Louis, MO). The library, on average, contained inserts of 120 kb with sizes ranging between 50 kb to 240 kb. A large genomic DNA fragment that contained VEGF promoter region was obtained. Southern blot analysis was performed to map the VEGF promoter region. A unique HindIII restriction site was mapped approximately 7.8 kb upstream of the ATG translational start codon of the VEGF gene. The sequences between HindIII and ATG translational start codon are inserted into the polylinker of pTKLR-Vn vector to finish the construction of targeting vector that contains VEGF-LucR transgene (Figure 3A).

#### 15 B. pTKLG-Fos/VEGFR2

A mouse VEGFR2 genomic DNA sequence of 1079 bp which contains a partial VEGFR2 promoter region was published previously (Ronicke, et al., (1996) *Cir. Res.* 79:277-285). Accordingly, primers were designed to amplify a 0.45 kb (KF1-KR1A; Table 1) and a 0.58 kb fragment (KF2-KR2; Table 1). It was confirmed that each pair of primers can amplify the predicted product using mouse129SvJ genomic DNA as

template. DNA sequences for these primers are shown in Table 1 above and PCR amplification conditions are shown in Figure 6. These primers were used for PCR screening of mouse 129/SvJ genomic DNA BAC library. From the screening, a BAC clone that contained a large genomic DNA fragment was obtained. Based on the published restriction map of VEGFR2, a 4.5 kb HindIII-XbaI fragment that covers the VEGFR2 promoter region was subcloned from the VEGFR2 BAC clone into the pBluescriptSK vector (Stratagene, La Jolla, CA) that was linearized with HindIII and XbaI. The VEGFR2 promoter sequences of 4.1 kb, spanning from a HindIII site to the ATG translational start codon, are subcloned into the polylinker of pTKLG-Fos vector to construct the targeting vector that contains VEGFR2-LucYG transgene (Figure 5A).

### C. pTKLG-Fos/Tie2

Mouse Tie2 genomic DNA sequence of 477 bp which contain a partial Tie2 promoter region has been published previously (Fadel et al (1998) *Biochem. J.* 330(Pt. 1):335-343). Accordingly, primers were designed to amplify a 0.45 kb (TF3-TR1; Table 1) and a 0.47 kb fragment (TF2-TR1; Table 1). It was confirmed that each pair of primers amplified the predicted product using mouse129SvJ genomic DNA as template. DNA sequences for these primers are shown in Table 1 above. PCR amplification conditions are shown in Figure 6. These primers are used for PCR screening of mouse 129SvJ genomic DNA BAC library. From the screening, a BAC clone containing a large genomic DNA fragment of the Tie2 promoter region was obtained. Based on the published Tie2 genomic DNA restriction map (see, Dumont et al., *supra*), a 10.5 kb Asp718-EcoRV fragment containing the Tie2 promoter region was subcloned ifrom the Tie2 BAC clone into the pSK vector that was linearized with Asp718 and EcoRV. The Tie2 promoter sequences of about 6.8 kb, spanning from the Asp718 site to the ATG translational start codon is subcloned into the polylinker of pTKLG-Fos vector to construct the Tie2-LucYG targeting vector (Figure 5B).

**Example 4****Generation of Transgenic Mice Carrying the Constructs of the Present Invention**

5       **A. General Procedure:** Figure 7 depicts a generalized description of generation of transgenic mice using the targeted transgenic vectors described in Example 3. Details regarding embryonic stem (ES) cell culture, transfection, blastocyst injection and implantation to a pseudopregnant foster are described, for example, in Hogan et al (1994) "Manipulating the Mouse Embryo, A Laboratory Manual. Second Edition", Cold Spring  
10 Harbour Laboratory Press.

After construction the targeted transgenic construct are transfected into C57BL/6 embryonic stem (ES) cells. (Genome System Inc., Genome Systems, Inc., St. Louis, MO) through electroporation. The antibiotic G418 is used to select for cells in which the DNA construct containing the Neo gene is integrated, either randomly or by homologous  
15 recombination. The nucleoside analog gancyclovir is converted by TK to a cytotoxic derivative. DNA that has integrated by homologous recombination lose the TK gene and are resistant to the drug, whereas cells that have incorporated the DNA randomly are likely to retain the TK gene. Thus, cells containing random integrations into a chromosomal location that allows the expression of the TK gene are killed. The G418  
20 and gancyclovir resistant clones are then be screened by PCR and Southern blot analysis and those that have homologous DNA recombination is used for FVB/N blastocyst injection (Genome System, Inc.). Between 4-16 blastocysts are transferred to the uterus of a pseudopregnant foster mother. The pups are typically born 17 days after the transfer. Either random bred mice or F1 hybrid mice make suitable recipients. Females of certain  
25 random-bred stocks (*e.g.*, CD1 mice, from Charles River Laboratories) have very large ampullae, which makes oviduct transfer easier. These mice also generally make good mothers. Alternatively, F1 hybrid females (*e.g.*, B6 x CBA F1) can be used as recipients. Although their ampullae are smaller, make exceptionally good mothers, rearing litters as small as two pups. See, for example, Hogan et al. (1994), *supra*.

**B. Screening for homologous DNA recombination positive ES cells**

- 1) **pTKLG-Fos/VEGFR2:** Analysis of homologous DNA recombination between pTKLG-Fos/VEGFR2 targeting vector and the FosB gene is carried out using Southern blot analysis as shown in Figure 8. Genomic DNA prepared from G418 resistant ES  
5 cells is digested with PvuII and probed with probe A to confirm the 5' end DNA recombination. PvuII digestion of DNA bearing homologous recombination reveals two separate bands of 8.2 and 4.0 kb, whereas digestion of DNA from homologous recombination negative clones reveals only the 8.2 kb band. The 3' end of DNA recombination is tested by hybridizing NotI digested DNA with probe B. NotI digestion  
10 of DNA bearing homologous recombination will reveal two separate bands of >8.2kb and 5.0 kb, whereas digestion of DNA from homologous recombination negative clones will only reveal the >8.2 kb band. Once homologous DNA recombination is confirmed, positive clones is selected for FVB/N blastocyst injection.
- 2) **pTKLG-Fos/Tie2:** Analysis of homologous DNA recombination between pTKLG-Fos/Tie2 targeting vector and the FosB gene is analyzed by Southern blot in a similar manner as described above for pTKLG-Fos/VEGFR2. Once homologous DNA recombination is confirmed, positive clones are selected for FVB/N blastocyst injection.
- 3) **PTKLR-Vn/VEGF:** Analysis of homologous DNA recombination between pTKLR-Vn/VEGF targeting vector and the vitronectin gene is analyzed by PCR. DNA  
20 primers designed according to the predicted homologous recombination, are listed in Table 2.

**Table 2**  
 PCR primers for analysis of homologous DNA recombination between pTKLR-  
 Vn/VEGF targeting vector and the vitronectin gene

5

<b>5'end primers</b>		
<b>F51</b>	5'- CCCAGTGTCTCTGATTTAGGGAGAGCACCTGAG -3'	SEQ ID NO:30
<b>R51</b>	5'- CCAGACTGCCTTGGGAAAAGCGCCTC -3'	SEQ ID NO:31
<b>F52</b>	5'- CAGTGAGAGTCTTCTCTGTCCCTCAATCGGTTCTG -3'	SEQ ID NO:32
<b>R52</b>	5'- TGGATGTGGAATGTGTGCGAGGCCAG -3'	SEQ ID NO:33
<b>3'end primers</b>		
<b>F31</b>	5'- AATCAAAGAGGCGAACTGTGTGTGAGAGGTCC -3'	SEQ ID NO:34
<b>R31</b>	5'- CGGCTCCCCAAAATGTGGAAGCAAGC -3'	SEQ ID NO:35
<b>F32</b>	5'- GAATCCATCTTGCTCCAACACCCCAACATC -3'	SEQ ID NO:36
<b>R32</b>	5'- CGCCTCCTCTCCCCAGTCTCCCCTTG -3'	SEQ ID NO:37

Primers F51-R51 and F52-R52 amplify a 1799 bp and a 1841 bp DNA fragment respectively from the 5'end of the transgene that is integrated into the vitronectin site through homologous DNA recombination, whereas primers F31-R31 and F32-R32  
 10 amplify a 3549 bp and a 3428 bp DNA fragment respectively from the 3'end of the transgene that is integrated into the vitronectin site through homologous DNA recombination. Clones that allow successful amplification of both the 5'end and 3'end of the integrated transgene are selected for FVB/N blastocyst injection.

15

### C. Analysis of chimeric mice

The pups developed from injected blastocysts contain chimeras, as can be identified by their agouti coat color when an ES cell derived from a mouse having a dark coat color (e.g., C57BL/6) is injected into the blastocyst of a light coat color animal (e.g., FVB/N, genotype B/B). DNA analysis (e.g., Southern blotting, PCR) is conducted to  
 20 further confirm the presence of the transgene in these pups as described above in Section B. These animals may be obtained commercially, for example from The Jackson Laboratory, Bar Harbor, MN.

#### D. Generating targeted transgenic C57BL/6 mice with white coat color

Breeding of the chimeric mice generates homozygous targeted transgenic mice, as depicted in Figure 9. The targeted mice are used to monitor gene expression through the measurement of luciferase mediated light emission from the mice. In a preferred embodiment, the targeted mouse has a light coat color (e.g., white coat color), because the black colored coat (an example of a dark coat color) of C57BL/6 mice can absorb light emitted from the body and may interfere the sensitivity of the bioluminescence assay. An inbred mouse strain C57BL/6-Tyr C2j/+ strain (Jackson Laboratory, Bar Harbor, MN) is available for this purpose. This strain of mice have white color coat, yet they still have the same genetic background as C57BL/6 mice except that the gene responsible for the black coat color is mutated. Unfortunately, C57BL/6-Tyr C2j/+ ES cells are not currently available. Therefore, the designed breeding program illustrated in Figure 9 is aimed to generate mice that are homozygous for the target transgene and have white coat color. C57BL/6 ES cells are prepared as described above and introduced into a suitable blastocyst (e.g., from the FVB/N strain of mice). The blastocysts are implanted into a foster mother. Chimeric mice are shown in Figure 9 as white animals with black and green patches. Chimeric animals are bred with C57BL/6-Tyr C2j/+ mice to create F1 hybrids. Subsequent breeding of the F1 hybrids generates several type of mice, including the one that is homozygous for the target transgene and has a white coat color (shown in Figure 9 as b/b; L/L), which is used for *in vivo* gene regulation monitoring.

A C57BL/6 mouse and a C57BL/6-Tyr C2j/+ mouse are considered to be substantially isogenic. Accordingly, the method of the present invention exemplified in Figure 9 provides a means for generating breeding groups of substantially isogenic mice in a selected genetic background carrying at least one transgene of interest.

#### E. Dual luciferase targeted transgenic mice

As described above, two targeting vectors are generated. PTKLR-Vn carries a red luciferase gene and is targeted into vitronectin locus. PTKLG-Fos carries a yellow-green luciferase gene and is targeted into FosB locus. A number of promoters, including VEGF promoter, VEGFR2 promoter, and Tie2 promoter are cloned into these vectors, as described above. Subsequently three type of targeted transgenic mice are generated.

VEGF mice carry VEGF promoter-red luciferase transgene (VEGF-LucR) integrated into vitronectin locus. VEGFR2 mice carry VEGFR2 promoter-yellow-green luciferase (VEGFR2-LucYG) transgene integrated into FosB locus. Tie2 mice carry Tie2 promoter-yellow-green luciferase (Tie2-LucYG) transgene integrated into FosB locus. Through a  
5 breeding program illustrated in Figure 10, dual luciferase targeted transgenic mice are produced, carrying both of the VEGF-LucR and the VEGFR2-LucYG transgenes. The degradation of luciferin by yellow-green luciferase and red luciferase generates lights that emit at 540 nM and 610 nM respectively. These wavelengths of light are measured individually using a photo-counting camera (intensified CCD). Therefore, both VEGF  
10 expression and VEGFR2 expression, for example, can then be monitored in the same mouse at the same time.

As is apparent to one of skill in the art, various modification and variations of the above embodiments can be made without departing from the spirit and scope of this  
15 invention. These modifications and variations are within the scope of this invention.



CLAIMS

What is claimed is:

1. A vector for use in generating a transgenic non-human mammal, said mammal having at least one single-copy, non-essential gene in its genome, said vector comprising

(a) a first selectable marker and a reporter expression cassette, said reporter expression cassette comprising a transcriptional promoter element operably linked to a light-generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, said targeting polynucleotide sequences flanking (a), wherein (i) the length of the targeting polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) said transcriptional promoter element is heterologous to the single-copy, non-essential gene.

2. The vector of claim 1, wherein said first selectable marker provides a positive selection.

3. The vector of claim 2, wherein said first selectable marker is selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

4. The vector of claim 3, wherein the first selectable marker is neomycin phosphotransferase II.

5. The vector of any of claims 1-4, further comprising a second selectable marker, wherein at least one target polynucleotide sequence is located between said second selectable marker and (a).

6. The vector of claim 5, wherein said second selectable marker provides a negative selection.

7. The vector of claim 6, wherein said second selectable marker is selected from the group consisting of adenosine deaminase, thymidine kinase, and dihydrofolate reductase.

8. The vector of any of claims 1-7, wherein the transcriptional promoter element is selected from the group consisting of an inducible promoter, a repressible promoter, and a constitutive promoter.

9. The vector of claim 8, wherein the transcriptional promoter element is selected from the group consisting of VEGF, VEGFR, and TIE2.

10. The vector of any of claims 1-9, wherein the sequences encoding the light-generating protein are obtained from either procaryotic or eucaryotic sources.

11. The vector of claim 10, wherein the light generating protein is a luciferase.

12. The vector of any of claims 1-11, wherein said reporter expression cassette further comprises other control elements.

13. The vector of claim 12, wherein said control elements are selected from the group consisting of transcription enhancer elements, transcription termination signals, polyadenylation sequences, sequences for optimization of initiation of translation, and translation termination sequences.

14. The vector of any of claims 1-11, wherein said vector is circular.

15. The vector of claim 14, wherein said vector contains at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: targeting polynucleotide sequence - (a) - targeting polynucleotide sequences.

16. The vector of claim 5, wherein said vector is circular.

17. The vector of claim 16, wherein said vector contains at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: target polynucleotide sequence - (a) - targeting polynucleotide sequences - (second selectable marker).

18. The vector of any of claims 1-17, wherein the coding sequences of the reporter expression cassette comprise codons that are optimal for expression in a host system into which the expression cassette is to be introduced.

19. The vector of claim 18, wherein said mammal is a rodent.

20. The vector of claim 19, wherein said mammal is a mouse.

21. The vector of any of claims 1-20, wherein said targeting polynucleotide sequences from single-copy, non-essential genes are selected from the group consisting of vitronectin, *fosB*, and galactin 3.

22. A method of producing a transgenic, non-human mammal, said mammal having at least one single-copy, non-essential gene in its genome, comprising  
transfecting an embryonic stem cell of said mammal with a linear vector comprising

(a) a first selectable marker and a reporter expression cassette, said reporter expression cassette comprising a transcriptional promoter element operably linked to a light-generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, said targeting polynucleotide sequences flanking (a), wherein (i) the length of the polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) said transcriptional promoter element is heterologous to the single-copy, non-essential gene;

selecting embryonic stem cells which each have said first selectable marker and reporter expression cassette integrated into its genome;  
injecting said embryonic stem cells into a host embryo,  
implanting said embryo in a foster mother,  
maintaining said foster mother under conditions which allow production of an offspring that is a transgenic, non-human mammal carrying said reporter expression cassette.

23. The method of claim 22, wherein said offspring is capable of germline transmission of said reporter expression cassette.

24. The method of claim 23 further comprising breeding said offspring with a mammal, wherein the mammal is substantially isogenic with the embryonic stem cells, wherein said breeding yields transgenic F1 offspring carrying said reporter cassette.

25. The method of claim 24, further comprising breeding a first F1 offspring carrying said reporter cassette with a second F1 offspring carrying said reporter cassette, wherein said breeding yields transgenic F2 offspring carrying said reporter cassette.

26. The method of any of claims 22-25, wherein said mammal is a mouse.

27. The method of claim 26, wherein said embryonic stem cells are derived from a mouse having a dark coat color.

28. The method of claim 27, wherein said mammal substantially isogenic with the embryonic stem cells has a light coat color.

29. The method of claim 28, wherein said F2 offspring carrying said reporter cassette has a light coat color.

30. The method of claim 29, wherein said embryonic stem cells are derived from a C57BL/6 mouse having a dark coat color, and said mammal substantially

isogenic with the embryonic stem cells is a C57BL/6-Tyr C2j/+ mouse having a light coat color.

31. The method of any of claims 22-30, wherein the light generating protein is a luciferase.

32. A transgenic, non-human mammal, comprising at least one single-copy, non-essential gene in its genome, wherein (i) at least a portion of at least one single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the gene, such that the gene cannot produce a functional gene product, and (ii) said polynucleotide sequences comprise a first expression cassette which has been introduced into said mammal or an ancestor of said mammal, at an embryonic stage, said first expression cassette comprising

a first selectable marker,

a first transcriptional promoter element heterologous to the gene, and

light-generating protein coding sequences, wherein said light-generating protein coding sequences are operably linked to said promoter element.

33. The transgenic, non-human mammal of claim 32, wherein the single-copy, non-essential gene is selected from the group consisting of vitronectin, *fosB*, and galactin 3.

34. The transgenic, non-human mammal of any of claims 32-33, wherein the first selectable marker is selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

35. The transgenic, non-human mammal of any of claims 32-34, wherein the first transcriptional promoter element is selected from the group consisting of an inducible promoter, a repressible promoter, and a constitutive promoter.

36. The transgenic, non-human mammal of claim 35, wherein the first transcriptional promoter element is selected from the group consisting of VEGF, VEGFR, and TIE2.

37. The transgenic, non-human mammal of any of claims 32-36, wherein said expression cassette further comprises other control elements.

38. The transgenic, non-human mammal of claim 37, wherein said control elements are selected from the group consisting of transcription enhancer elements, transcription termination signals, polyadenylation sequences, sequences for optimization of initiation of translation, and translation termination sequences.

39. The transgenic, non-human mammal of any of claims 32-38, wherein said non-human mammal comprises a second single-copy, non-essential gene in its genome, wherein (i) at least a portion of the second single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the second gene, such that the second gene cannot produce a functional gene product, and (ii) said polynucleotide sequences comprise a second expression cassette which has been introduced into said mammal or an ancestor of said mammal, at an embryonic stage, said first expression cassette comprising

- a second selectable marker,
- a second transcriptional promoter element heterologous to the second gene, and
- light-generating protein coding sequences, wherein said light-generating protein coding sequences are operably linked to said second promoter element.

40. The transgenic, non-human mammal of claim 39, wherein the first transcriptional promoter element in the first expression cassette is different from the second transcriptional promoter element in the second expression cassette.

41. The transgenic, non-human mammal of claim 40, wherein the light-generating protein in the first expression cassette can produce a different color of light relative to the light-generating protein in the second expression cassette.

42. The transgenic, non-human mammal of any of claims 32-41, said mammal being a rodent.

43. The transgenic, non-human mammal of 42, said rodent being a mouse.

44. The transgenic, non-human mammal of any of claims 32-43, wherein the sequences encoding the light-generating protein are obtained from either procaryotic or eucaryotic sources.

45. The transgenic, non-human mammal of claim 44 wherein the light-generating protein is a luciferase.





FIG. 1








FIG. 3A




FIG. 3B

1 TCCAGCCACC TTTTCTAC GTCGCCGCC TTCTAGTTA ACTTCATGT TAAAGAGCC TCCGCCGCC AGGTGTGTG CCCACAGAG GAGGCTGCT  
AGTGTGTG AAGAGAGT CAGGCCGCC AGGATCAAT TGAATACCA ATTCTTCCG ATGTGCCGCC TCCACACCA CGTGTCTTC CTGCCACGA

101 CCCACAGCC CCCAGTGTG CTGATTAGG GAGAGCACT GAGGCCATG AGAGTCTCT CTGTCCCTA ATCCGTTCG AATTCGCCA CTGCCCTCC  
CGTGTGTG CGGTACAGA GACTAATCC CTCTGTGGA CTGGGTGAC TCTCAGAGA GACAGGAGT TACCCAGAC TTAAAGGCT GAGCCGAGG

201 TTATCCAGG GACAGGCTG CCCAGCTAT TCAGAGAGT AGTCTTAAC TGTAGCCAA CAGACTTTT ATTGGCTGG GAGAAAGAG TAAAGCTCT  
AATAGTCCC CTGTCCGAC GGTGGGATA AGTCTGTCA TCAGATTTG AGCATCGTT GTCTGAJAA TAACCGGACC CTCTTCTCT ACTCCAGGA

301 GAGGCTCAG CGAGTGGCT CTGATTCCT CTCTCCAGG GTCCGCCAG CCAGCCATA CTGAGCAAT GAGCGTGTG AGGAGGTT CACAGAGTC  
CTTGGATGG GTCACCCGA GACTAGGAT GAGAGTCTC GAGGAGTCC GGTGGGTCT GACTGTGAC CTGCCACCA TCCCTCTTA GTGTCTCAG

401 ACTGCCCGG TCTCAGGTT GACTGGTAG TATTGTCTG AAGAAAGAA TGAJAAAGG GTATGTGAG ATTCTGGCT ATCTGTCCA CTGTGCCAA  
TGAGCGCCC AAGATTCCA CTGAGCATC ATAAACAGAC TTCTTTCTT ACCTTTTTC CATACACTC TAAGACGAC TAGGACAGT GAGCGGTT

501 GAGGATAAA GCTTTTCTC CAGAGGGA AGTGAACATC CAGCAAGAG ATATGTAC CATCTACAG CTGTGTTCG CAGCCAGGA CCAAGAGCT  
CTTCTATT CTGAJAAAG GTCTTCCCT TCACTTGTG GAGAGTCTC GTGTGTGCT GAGACAGT GAGACAGT GAGACAGT GAGACAGT

601 CAGGCAAGC CTACCCAAA CAGCTTAGG GTCTGATTC CTCTCTTC GGTGCCAC CTCCAGAA GAAATAGAG CTCTGAATG GCTGCCAGT GCAAGTACA  
GTCTGTTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC

701 AGGAGTCCA TATCATATC ATAGTTTGT TAGGTCTCA GGCCTCTCT CTGCTGGAG AACAAAGAG ACCAGATGA ACCTGTGAA CAGCGGAGT  
TGGTCAAT ATATATATC TATCAGAC ATCCAGAGT CCGGTGAGG GAGGAGCTC TTCTTCTCT TGTCTTACT TGTCTTACT TGTCTTACT

801 TGGAGCTCT GCTGGGTCT TCCGCCGCC CTGCGGTGA AGATAGGCT TTTCGGCTC TACGCTTGA CTCTGTCTT TTGGCTTGG CAGGCTGGA  
AGTGTGTG AGGATGAG AGGATGAG AGGATGAG AGGATGAG AGGATGAG AGGATGAG AGGATGAG AGGATGAG AGGATGAG

901 TAAAGAGCA GTAGGTAGA TCCGCCGCC CATAGAGG TCCACTTTC CTGCGACAC ATGCGATTC GCGGTGATG ATTGGGTTT TGTGCCCA  
ATCTCTGT CACTGATCT AGCGCGGCC GTATGTGCT AGGTGAAGG GAGCGTGTG TACGCTTGA GCGGCTTACT TAAGCGAGG AGCGCGAGT

1001 TCTGAGAGC GAGAGGCTT AATGACTTG GCAGATCTG GCTTTGATT CTCCAGAGG GTGTCTCTC TATCTATTA TGTATCTTA TGTATCTTA  
AGACATCT CTTGCCCA TTAGGTGAG GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC GTCTGATTC

1101 TATCTATTA TGTATCTTA TATCTATTA TATCTATTA TATCTATTA TATCTATTA TATCTATTA TATCTATTA TATCTATTA TATCTATTA  
ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA ATAGTATTA

1201 CCTATTATT TTTTGTTC TTTTCTTGA AACAGATCT TACGAGCTC CTATGCTGG TTTCGACTC ACTATGAGC CATAGCTGC CTCTTACTC  
GATTAJAAAG ACAAJAAAG AAAGAAACT TTGTCTTGA ATCTGTGAG GATAGGACC AAAGTGTAG TGTACTTCT GTATTGAGG GATATTGAG

1301 ACAAGATCC ACTTGGCT GTCTCTGAT GCTGGATTA AAGCATGT CCACTACAC CAGCTCAGT AGGAGCTTA GAGCATATT GCTATGCTT  
TCTTGTGAG GTAGGAGCA CAGAGCTCA GAGGCTTAT TTCTGATAC GTGTATGAG GTGTATGAG GTGTATGAG GTGTATGAG GTGTATGAG

1401 GCTTAGACA CCAACTCAG TCCGCCGCC CCAGGCTCC TGTCTAGG TTTTCCCAT CTTCTCTCA CTGTATCTT TGAATCTCT CCGCATGCC  
CGGATCTCT GTTGTGATC AGGCGTGGG GTTCCAGAGG ACAGATCTC AAAGAGGTA GAGAGAGGT GACATAGGA ACTTAGAGC GGGTGGCT

1501 AACCCCTAG CCGCAGCCC CTCTCTCTC TGTGTAGG AAGTCCAG GTATGGATC CAAATAGAG CAGGCTCAT CCGCAGAGG TCAACAGAG  
TTGGGAGTC GCGGTGGG GTTCTCTCT TGTGTAGG AAGTCCAG GTATGGATC CAAATAGAG CAGGCTCAT CCGCAGAGG TCAACAGAG

1601 CAAAGCTAG CAGAGCTAA CAGCTCTGA TGTATGAT CAGATTTCA GCGCCCTCC CTGAGAGCC CCACTATAC AGCCGATTT CAGAGAGG  
GTTCAGATC GTTCTGTTC GTTCTGTTC GTTCTGTTC GTTCTGTTC GTTCTGTTC GTTCTGTTC GTTCTGTTC GTTCTGTTC GTTCTGTTC

1701 AAGCAGGCT TGTCTGCT CCACTAGCA GATCTGCT CAGAGAGG GTTGTAAAT GTTCTCCAG CTGTCCGCT GAGCAGGC AAGTCTCA  
TTGGTCCA AGAGAGGA GTATGTCT CCAAGAGG GTTCTCTCT CAAAGCTTTA CAAAGAGG GTTCTCTCT CAAAGAGG GTTCTCTCT

1801 AACAGGCT CAGAGAGCT GCTTCCCTC TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT  
TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT TGTCTGCT

1901 ATCAGAGAG ATCTAGACA GCGAGAGG ATCTGTGA TAGAGAGG ACTCCAGAC CTGCGGCTT CTCTCCACC CTGATAGCC TTGAGAGT  
TAGTCTCT TAGATGCT CCGTCTCTC TTAGAGCTT TAGAGAGG ATCTGTGCT TTAGAGCTT TAGAGAGG ATCTGTGCT TTAGAGCTT TAGAGAGG

2001 AGAGCTTTC CCGGCACTG TAGAGGCTG CAGAGAGG GAGGCTTCA CTACATCTT CTGATAGCT GAGGAGTCT TGTAGAGT TGTAGAGT  
TGTAGAGG GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT

2101 ACCAGGCTC AGGAGGCT CCAAGAGG CCACTATGA CTAGCTTGA AGACAGAT CAGAGGCTG AGGAGTCT GAGGAGTCT GAGGAGTCT  
TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG

2201 TATATCTTA ACTGCTCTC ACTGCTCTC GAGTGGCT ACTGCTTCT CTTCAGACT CCGGAGCAG TCCATGCTC CTTTACTCT GCTTACTCT  
ATATAGAT TTAGAGAG CCACTGCTC TTAGAGAG CCACTGCTC TTAGAGAG CCACTGCTC TTAGAGAG CCACTGCTC TTAGAGAG CCACTGCTC

2301 TAGGCTGCT ACTTGAACA AGTAGGCTT CCGGAGAG TTAGAGAG TTAGAGAG TTAGAGAG TTAGAGAG TTAGAGAG TTAGAGAG  
ATCCAGCA TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT

2401 TTGGGTTAC CAGGCTCAG TCCGCTTCA TCTAGCTAT AGCAGTACT CCACTAGCA GCGGAACTG TGTAGAGG AGATAGGCT AGGAGAGT  
AAGCCATG GTCCGAGT AGGAGAGT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT TTAGAGCT

2501 CCAAGGCTC CAGAGCTCT CATAGATCA CCAAGAGG CAAAGAGG CCAATCTCT AGATCCCTG AGGCTCAA GCGCTTCA CTGCAGTT  
GCTGCGGT GTCTGAGG GTATCTCAT CCAAGCTCT CAAAGAGG GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT GTTCTGCT

2601 CTGCTCTG AAGCTAGG GTCCCTGAT CAGTGTCT GCGGCTTAG ATCTCTCT GTTCTCAG TTAGAGGCT GCGGCTCTG GCTTCTCT  
GAGGAGCT TTGTGCTC CAGGAGCT GTCCAGAG GTCCAGAG GTCCAGAG GTCCAGAG GTCCAGAG GTCCAGAG GTCCAGAG GTCCAGAG

2701 AGGATCTAG AGGCTGCT CTTTAGAGT GCGTCCCT CAGAGCTTA GGTCAAGG TGTAGAGG TGTAGAGG TGTAGAGG TGTAGAGG  
TCTAGATC CCAAGCTC CCAAGCTC CCAAGCTC CCAAGCTC CCAAGCTC CCAAGCTC CCAAGCTC CCAAGCTC CCAAGCTC

2801 TTCTTGGCT CTTCACTA GTCAAGCTC CAATATCAT CTTCTGCT AGTGAACAG TCCCGGCTC TTAGAGGCT CAGAGGCTC AGGAGAGT  
AAGAGAGCA GAGATGCT CAGATGCT GTATATTA GTATATTA GTATATTA GTATATTA GTATATTA GTATATTA GTATATTA GTATATTA

2901 GTAGGCTCT GAGGAGGCT CCAAGGCTC CCAAGGCTC TGTAGCTA CTTTGGCT TCCATCTCT CAGAGGCTC CAGAGGCTC CAGAGGCTC  
CAGTCCAG CTTCCCTCT GGTCCGCTG GTTGTGCT AGATATTA GAGAGGCT AGGAGAGG GTATAGCT CAGATGCT

FIG. 3C-1

3001 AGTCAAAAGC TGTTCACACT GACACTTCTT GCTGCGCATG AAACCCCTGAG TGCAGCCGCC CTTCGATGAC TGTATGGGAG GGAATATGAG GTTTACAGCC  
 TCACTTTTGG AGCAGTGTGA CTGTGAGAA GACCCGCTAC TTTCGGACTC AGCTCCCLGG GAAGCTACTG AGATACCCCTC CCTATATAGTC CAATATGTCG  
 3101 CAATCTAGGG CAGCTGDCCA ACCTGACATT CCTAGGCTAC CCACCAATCC CCTCCACAC CTTCGCTCAGC CAGAGAAACC CAGTCCACCA GCGCTAGGAT  
 GTTAGATGCC GTGACGGGT TGGAGGTGAA GGGATCCATG GGTGCTTAGG GAGGGGTGTG GAACCACTCG GTCTCTTTGG GTACCGTGTG CCGGATCATA  
 3201 GAAAAAGGCC CTCAGGGGTG CCACTGCGAG CTTCTAGGCC AGGGCTTTGG CAGCTGCGCC GCGAGCTTC TGGATCTCG CTCTCTGCG TGAJAAAAA  
 CTTTTTCCCG GAGTCCCGAC GGTACCTTCC GGAGATCGGG TCCCGGAAAC GTTCGACCGG CCGCTCGAAG ACCTTAGAGC GACAGAGCGG ACTTTTTCT  
 3301 AGCAGACTGA AGAAGAGTTC CTAGTTCCTT GGTTTCTTGC GGTATATTTG CTCAGCTCTT GCGCCAGGCC CATTGCGCTC CTCGAAACAC AGCTGACCA  
 TCTCTGACT TCTTCTGAG GATCAGGGA COCAAGACCG GGAATTAAC GAGTAGGAGA CCGGTTCGGG GTACCGGAG GAGGTTTGTG TCGAGTGTGT  
 3401 AAGGCTACA TTCCAGAC CCAGCCCA GAGAGCTGG GAAACAGAA ACCCTCCCA AGACCAAGT CAGTAGGGTC AGGGGAGGA GGGATACAC  
 TTCCAGTGT AAGGCTCTG GGTTCGGGT CCTCTCGAC CTTTCTCTTT TGGAGCGGT TCTGTTTCA GTCTATCCAG TCCCGTCTT CCTATTTGT  
 3501 GCTTAGCTTA GCTGGGAGG TGGAAAGAG CATGTGTGT CACCCCTGA GCGAGTCCCG TTAATCTCC TGAAGCTTAC TTTTATAA GTGGAGCAT  
 GGAATCGAAT GAGTCCCTTC ACCTTCTTC GTACACACA GTGGAGACT CCGTCAAGGC ANTAGAGGG ACTCGGATG AAAATATAT CACCTGTGTA  
 3601 GGTGCTTGC CTATCAAGT GTTAGAGAT TCGTAGCT AGAAGACGA AAAGCTTTG TCGCTGAGT AGCTTCCAC TCATTCGAT AAGCGTAT  
 CCAAGGAAG GAGTAGTCA CACTCTCTA AGGACTGGA TCTTGTCTT TTTCAGAGC AGGACCTCA TCGAAGGTG AGTAGGGTA TTGCGCAATA  
 3701 GAAATTAAGT TTTAGTACG CTAGTCTT GTCCATCTT ACCCGCCCT TCGAATCTG ATTTTGGGG CAGAGAGGG GGTTCGGGA GAGCTGCA  
 GGAATGAC AAATAGTCC GATCAGGA GAGGTAGGA TGGGGGGGA TCTTAGAGC TAAJAAACCC GTTCTTCCCG GTTCTGCGG GAGCTGCA  
 3801 GCACTTTGG GAGGTTTTC TTTTCTTC ABAAGAGAC AAAGCTTCT TTTCTCTG TTTCAGGA AGAGCGGAG AGGACAGGA GAGATCGAC  
 CTTGAAAGCC CCTCCAAAG AAAGAGAGG TATTTTCTG TTTCAGGA AGAGCGGAG AGGACAGGA GAGATCGAC CCAAGATGTC GTATCTCTCA  
 3901 AGTGGTCTG AGTCTATCT TCTTCTTA TTTTCTTG ATTTATAT TTTATGTTT GTGTATAGT GTCTGCTAC ATGTGACT GTGACCA  
 TCACCCATC TCAAGTAGA AGAAGAAAT AAAAATAAT TAAATATAA AATACAAA CACTATTTA CAGAGATG TACAGTGA GTGAGTGT  
 4001 TCGATCTCT GTCTCTAGG AGGTAGAG AGGCTTTGA ATACCTGGA ACTGAGTTT TGAACATTA TGAAGTGG TGTGATGCT GAGATCTAA  
 AGTACAGAA CAGAGTACC TCGATCTT TCCGAAAT TATGGACT TATGCTCAA ACTTGTCAAT ACTGAGGCG ACAGCTAGGA CTCTAGTTT  
 4101 CCCAGTCTT CTGTAGAC AGTACTCTT AAAGCTGAG CCATCTTCC AGTCCAGAG CCGATCTCT AGGCTTTAC TATTCATG ATCCTGCGG  
 GGTTCAGGA GACTTCTT TCGATGAA TTTCGACTC GTTAGAAGG TCGAGGTCTC GGTTAGGAC TCGAAGGTG ATTAGGTAC TAGAGCGCC  
 4201 GACCACTCT GGCACACTT CATGACTC ATTTATTTA AAAAATAAT GACTCATG GGCATCTT CTAGACTAC ATCTAAGT GATTTCTCT  
 GTGTGGAC GGTGTGGA GTTAGTGA TATTTTCTA TTTTCTTA CCGTAGTAC CCGTAGTAA GATCTAGT TATGATCAC CTAAGAGGA  
 4301 ABAAGAAAT GCTTACTGG GTAGATGCC AGTTTGGG CCAATTOCA AGCACTGCA CACTTCTGA GCGCTCTCT TTTCTTCT GTATCAGG  
 TATTTCTTA CAGTGAAC CATCTACGG TCCAAAGCC GTTAGAGT TCGTAGCCT GTAGAGACT CCGGAGGCA AAAGACAGA CATTAGTCT  
 4401 GCGAGCTGC CTTCGCTGC TCTTCTCT GAGCGGAGT AGTCTAGG CCAAAATGA ACATTAAT TACTCTCTA CAGAGCGT AAGCTTAT  
 CCGTCCAGC GAAGACAG AGAGAGTA CCGGTCTA TCGAGTCC CTTTACTT TGTATTGA ANTAGGGAT GTCTCCGAC TTCTGATTCA  
 4501 GGAAGCGCC ATTAAGGCC TTAAGAAAT TCAACTGGA TCTTTTACC ATCCGAGGG GAGTGGAT CATGTAGCA GCTTCTTCC ACATTTTGG  
 CCTTTGCGG TAAATTCGG AATTTCTAG AGTTGAGCT AAGAAATGG TAGGCTGCC CTGCACTAT GTACATCGT CCAAGAGAG TGTAAAGCC  
 4601 GAGCGAGCG AGCGTAGGA AATGAGAG AGCTCTTAC AGGCTTTCT ACAGCATCTT GCACAGCAC AAGGGAGAC TGGGAGAG AGCGGAGCG  
 CTGCGCTGC TCCCATCTT TACTTCTG TCGAAGAT TCGGAAAG TGTCTAGAA CGTGTGTG TTCCCTCTG ACCGCTCAG TCCCGTCTG  
 4701 AGGTGTGGC GTGCTGGAG ACCTGGGTA GCTTTGCGG TCGTTCGGG GCGAGCGCG TGAACCTAG AGCGGGGCG TCAATCTCT GACTCTGCT  
 TCCACAGCG CAGGAGCTC TCGACCCAT CGAAGCGCG AGCGAGCGG CCGCTGGGC ACTTTGGATC TCCGCGCGG AGTTTAGGA CTGAGAGAC  
 4801 CTCAGAGCG TCGTCTCTT TGAAGTCT AGCTCCGCT TCGTAGAT GAGCAGCGC TTTTCTCGG GCAGCGGCT CTCTAGCTC CCGCTCTG  
 GACTCTGCG ACCAGACA ACTGAGGA TCGAGGAG ACGAATCTA CCTCTCGG AAACAGAGC CTTGCGGCA GAGATGAGG GCGCAGAGC  
 4901 TCGAGCTTC TCTTCTCTT CATGCTCT CTAAGTCTT GATTCGCGA GCTGCGTCC TCTTCTCT TCTAGCTT TCGGAGCA CCTTACCG  
 AGGTACGAG AGAGAGGA GTAGCGGAG GATTCAGGA CTCAGGCT CAGCGGAG AGAAGAGGA AGATGTGAC ATCGGTCTT GGAATGCGC

FIG. 3C-2




FIG. 4A

1 GCAGCTGCCC AACCTTTGCC GATGCCCCGT CAAAGTATAT ACCGCTGTGT TAGCAGAGC TGAGAACTTT TAGCCAAAG CCGGCTGCTT AGCCGAAJGC  
CGTGGACCGG TTTCGAAAGG CAGCGGCCAC GTTTCATATA TGGGCGACCA ATGCTCTTCC ACTCTTGAAT ATCGGCTTTC GCGCGAGGGA TTGCGCTTTC

101 TAGGCAAGTA GGGGAGAGAA AAAAAAATA AATTTCCAGA GAAGCTTCCA GAGGCTGCTC CTCTTCCCTC TTCTTTCAAA AGGACTGCAA AGCCGCAATC  
ATCGCTTCAT CCGCTTCTTT TTCTTTTTTT TTCTAGGTCT CTTCGAAAGT TCGAGAGGAG AAGGAAAGTTT TCGTAGGTTT CAGGCGCTAG

201 ACCGCTGACC CAGCAGAGAT TAGGCTGCTC AACCCCGGTC ACCGCTGCTC CCGCTGCTCC GGAAGTTAAC GCGGAGACCG TCGTAGAGC CAGACCGGCT  
TGGAGAGTGG GTCTGTCTCA ATCCCGAGAC TTGGGCGCAG TCGGAGCGAG GCGGAGAGAG CCGTTCATTC CCGGCTGCTC AGCATTTTCG CAGTCCGCGA

301 GGAATCTGTC GTCTGACCGG GCGCAGCGAC AGCCCGGAGC CCGTCCGCGC GCGGCGCGCG TGAGCTGCGG GCAGCTTCTA TTTTGGAGG CCGAGCGCAC  
CGTTAGAGAG CAGACTGCGC CCGCTGCTTC GCGCAGCGAG TCGGCGCGAG CCGGCGCGCG ACTGCGAGCC GTTGCAGAT AAACCTTTCG GCGTCCGCTC

401 GTTCTGAGAG GAGCGGCGAG CCGTCCGCTTC TGATTTGGCT TCGGCGCGAG CTTCAGCCAA TGAGCTTTCG CTCTCTATTT GTAGAGGCTA GCGTCCGCTC  
CAAGGATTCG CCGGCGCGCT GCGCAGCGAC GCGCAGCGAG ACTGCGCTTC GATCTCTTTC GCGATCGGGA CTCTCTTTCG CCGCATGCTC TCGTAGGACT

501 TTGCTTTTTT TGCTTCTTTC TGCTTCTTTC GCGCAGCGCG GTCTCCAGGA GAGAGGCTAG GATCTCTTTC GCGATCGGGA CTCTCTTTCG CCGCATGCTC TCGTAGGACT  
AACGAAAGAC ACCGAGAGAG GCGCAGCGCG GTCTCCAGGA GAGAGGCTAG GATCTCTTTC GCGATCGGGA CTCTCTTTCG CCGCATGCTC TCGTAGGACT

601 TGTGTGAGAG TGCTTCTTTC TCAAGAGCTA GAGGCTTTTC GCTGAGTTTC AGGCGCTCTA AGGCGGAACT TCGGCTTTCG CCGCATGCTC TCGTAGGACT  
ACACAGCTTC ACCGAGAGAG GTATTTCTAT CTGCGAAAGC CTCTCCAGTC CTAGAGAGAG CCGTCCGCTC GCGATCGGGA CTCTCTTTCG CCGCATGCTC

701 ACCTGCTGCT GAGCTCTAGA CACTCAGTTC TTCCGAGGTC TTCAAAGACT AGATAGCTA GCGTCCGCTC AGGCGGAACT TCGGCTTTCG CCGCATGCTC TCGTAGGACT  
TGCAGAGGGA CTGAGAGGCT GTAGGCTCAG AAGGCTCCAC AATTTTCTGA TCTACTGAT CCGTCCGCTC GCGATCGGGA CTCTCTTTCG CCGCATGCTC

801 CTTCGCTTTC TTCCGAGGTC CTGATTTGCC AGGATTTCCG CCGTCCGCTC GCGGAGAGAG CCGTCCGCTC GATCTCTTTC ATTCGAGGAT CCGTCCGCTC TCGTAGGACT  
GAGGAGAGTC AAGGCTCCGA GAGCTCTAGA TCGGCTCAG TCGGCTCAG TCGGCTCAG CCGTCCGCTC GATCTCTTTC ATTCGAGGAT CCGTCCGCTC TCGTAGGACT

901 GCGGCGCGCG GCGGCTGCTC GAGCTCTTTC GCGGAGCGAG ATCTATGTC ACCGCTTTCG CTGAGAGGGA GTCTCTTTCG CAGACTCTCT AGAGGCTA  
CGCGGCGCGG CCGGCTCAG CTGAGAGGA CCGGCTGCTC TCGGCTCAG GAGGCTTTCG TCGGCTCAG GAGGCTTTCG CAGACTCTCT AGAGGCTA

1001 TGCTCTTTCG TTCTCTGAAA CCGTCTTTCG TCACTGAGAA GAGAGAGGGA CTGCGAGGTC AGGCTTTCG TCGGCTTTCG CAGACTCTCT AGAGGCTA  
ACGAGAGGTC AAGTACTTTC GAGGCTCAG AGTAGGCTTC CTCTCTTTCG GAGGCTTTCG TCGGCTTTCG CAGACTCTCT AGAGGCTA

1101 ACTGAGGATC CCGGCTGCGG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG  
TGAGGCTTTCG GCGGAGAGTC CCGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG GAGGCTTTCG

1201 TGTGTGAGAG GTTGGGCGGT TTGCTCTTTC TGTGTGTGTC AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGTCTTTCG CAGACTCTCT AGAGGCTA  
ACACAGGCTC CAGGCTTTCG AAGGCTTTCG ACACAGGCTC AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGTCTTTCG CAGACTCTCT AGAGGCTA

1301 CCGGCTTTCG ACAGGCTTTCG GTCTCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
CGTGGGCGGT TGTGAGGCTC CAGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

1401 GTTGAAGAGG ATCTCTTTCG AAGGCTTTCG TCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
CAAGCTTTCG TCGGAGAGG AATGCTTTCG AAGGCTTTCG TCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

1501 TGTGTGAGAG CAGGCTTTCG GTCTCTTTCG TCTCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
ACAGGAGTTC GTTGGGCGGT TCGGCTTTCG CAGGCTTTCG AAGGCTTTCG ACACAGGCTC AATGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

1601 GTTGAAGAGG CAGGCTTTCG GTCTCTTTCG TCTCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
CATGCTTTCG CAGGCTTTCG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

1701 CTCTCTTTCG CAGGCTTTCG GTCTCTTTCG TCTCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
GAGGAGAGG CAGGCTTTCG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

1801 GCGTGTGCTC CTACAGGCTC GCGGCGCGAA GCGGAGAGG TCGGCTTTCG ACCGCTTTCG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
CGGCTTTCG GAGGCTTTCG GCGGCGCGAA GCGGAGAGG CCGGCTTTCG ACCGCTTTCG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

1901 GCGGAGAGG CCGGCTTTCG TCTCTTTCG TCTCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
CGGCTTTCG GCGGCTTTCG GCGGCTTTCG TCTCTTTCG TCTCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2001 TCGGCTTTCG GAGGCTTTCG AGGCTTTCG AAGGCTTTCG TGTGTGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
ACGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2101 GAGGCTTTCG TGTGTGAGG AAGGCTTTCG GCGGCTTTCG CAGGCTTTCG TGTGTGAGG AAGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
CGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2201 AAGGCTTTCG AATTTTTCG CTGAGGCTC GCGGCTTTCG CAGGCTTTCG AAGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
TTCTCTTTCG TCGGCTTTCG GAGGCTTTCG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2301 TGTGTGAGG AAGGCTTTCG AAGGCTTTCG AAGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
TTCTCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG TCGGCTTTCG

2401 AAGGCTTTCG GTTGGGCGGT ACCGCTTTCG GCGGCTTTCG GAGGCTTTCG GAGGCTTTCG TCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
TTCTCTTTCG CAGGCTTTCG TCGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2501 CCGGCTTTCG GTTGGGCGGT ACCGCTTTCG GCGGCTTTCG GAGGCTTTCG GAGGCTTTCG TCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
GAGGCTTTCG GCGGCTTTCG GCGGCTTTCG ACCGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2601 CAGGCTTTCG TGTGTGAGG TCGGCTTTCG CAGGCTTTCG GAGGCTTTCG TGTGTGAGG AAGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
GAGGCTTTCG GCGGCTTTCG GCGGCTTTCG ACCGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2701 ATGCTTTCG CCGGCTTTCG TCTCTTTCG AAGGCTTTCG CAGGCTTTCG TGTGTGAGG AAGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
TGTGTGAGG GAGGCTTTCG TCGGCTTTCG GAGGCTTTCG TGTGTGAGG AAGGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2801 CCGGCTTTCG TTTTCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
GAGGCTTTCG GCGGCTTTCG GCGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

2901 CAGGCTTTCG GAGGCTTTCG GCGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
GAGGCTTTCG GCGGCTTTCG GCGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

3001 GCGGCTTTCG ACCGCTTTCG TTCTCTTTCG TTTTCTTTCG GTCTCTTTCG CAGGCTTTCG AACTTCTTTC CCGGCTTTCG AAGGCTTTCG GCGGCTTTCG  
CGGCTTTCG TCGGCTTTCG AAGGCTTTCG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

3101 CCGGCTTTCG GAGGCTTTCG CCGGCTTTCG GCGGCTTTCG GAGGCTTTCG GAGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG  
GAGGCTTTCG CCGGCTTTCG CCGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

3201 TGTGTGAGG AATGCTTTCG GTTGTGAGG TTTTCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
AAGGCTTTCG TTTTCTTTCG GCGGCTTTCG GCGGCTTTCG GAGGCTTTCG GAGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG

3301 TGTGTGAGG TTTTCTTTCG CCGGCTTTCG GTTGTGAGG TTTTCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
AATGCTTTCG TTTTCTTTCG GCGGCTTTCG GCGGCTTTCG GAGGCTTTCG GAGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG

3401 CCGGCTTTCG GCGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG  
GAGGCTTTCG CCGGCTTTCG AAGGCTTTCG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

3501 CCGGCTTTCG GCGGCTTTCG AAGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG CCGGCTTTCG  
GAGGCTTTCG GCGGCTTTCG GCGGCTTTCG TGTGTGAGG AATGCTTTCG GCGGCTTTCG GCGGAGAGG CCGGCTTTCG GCGGAGAGG

FIG. 4B-1

3601 CAGTTCGGAG CGAGCTCTTC AGCCCCCTGG ATCAGCACCA AGAATATATG AGTGTGGGAT TGGCGGGTGG AAGCTACTCT GTGTGGTCCG TGACCAACAA  
 GTCAACCTTC CTGTGACAGG TGGGGGAGCC TATGTCTGCT TCTTACATAC TCACACCCCA ACCCGCCACC TTGATGAGA CACACACAGG ACTGTGCTTT  
 3701 TTCTCTCTTC TCTGTCTCTC ATGACCTGCG CCTGCTGGGA TCGATAGGGA AACTGATCAG CTTGAGAGAG AAGAGCCAGA CCTGGATCCG GAGATCTCCG  
 AGACAGAGAG TACTGAGCGG GTACAGACCTT TGTACTAGTC TGTACTAGTC GAACCTCTCC TTTCCTGCTT CACTGATCAG CTCAGACCGC  
 3801 AGCTGCAAAA AGAGAGAGAA CGCCTGAGAT TTGTCTGCTT GCGCCACAAA CGCGGCTGCA AGATCCCTTA CGAAGAGAGG CGCGGGCCAG GCGGCTGCGC  
 TGACCTGTTT TCTCTCTCTT CGCGACCTCA ATCAGAGACCA CCGGCTGTTT GCGCGGAGCT TCTAGGGAT CTCTCTCCCG GCGCGGCTGC CGCGGAGCGG  
 3901 CGAGGTGAGA GATTTCGAGG GTTCACATTC CGCTAAGGAA CGCGCTTCG GTGTGCTGCT GCGCGGCGCT CCACACCGCC CACTGCGCTT CAGAGCAGC  
 GCTCCTCTCT CTAAAGCTTC CGAGTTTGAG GCGATTTCTT CCGGAGAGC GTGCGGAGC CGCGGGGAGA GTGTGTTGCG GCGAGCGGAA GTCTCTGCTG  
 4001 CGAGAGCCAC CGCCACACCT GACGCTCTCT CTCTTTACAC ACATGTAGAT TCAAGTCTCT GCGGACCGCT TCGCGCTTGT TAGCGCTTCT TACACTTCTT  
 GCTCTCGCTG GCGGCTTGA CTGCGGAGAA TGTCTACTTA AGTTCAGAG AGCTGCGGAG AGCGGCAACA ATCGGAGAGC ATGTGAGAGA  
 4101 CGTTTGTCTT CACTTCCCGG GAGGTCTCGG CGTTTCCCGG CGCGGACGCG ACCAGCGGCA GCGGAGCAGC GTCCGACCGG CTGAACCTCG CCTGCTTCTT  
 GCAACACAGA GTGAGACCGG CTGAGAGGCG GCAAGCGCGG GCGGCTTCCG TGTTCGCGCT CAGCTGTGCG CAGGCTGCGC GACTGTAGCG GAGGAGAGAA  
 4201 TCTCTCTGAA ACTCTTTAGA CAACAAAACT GTTTGTTTTG ATCTAAACCC CGTCTCTGCT TCTCTCTCTT TCTACTCTCT CTCTGCTCTC CTGTGCTGCG  
 AGAGAGATTT TGAGAAATCT GTTTGTTTTG TTTGTTTTGCG GTTCTCTCTT TCTCTCTCTT TCTACTCTCT CTCTGCTCTC CTGTGCTGCG GCGGCAACAA  
 4301 GTGTGAGAGG TTGACTCTCT CTGTCTGAGC AGCTGCGCGC TCTGCTCTCG GACTGAGAGG AAGGAGCTCG TTGTGTGTTT GTGTCTGTCT TCTGCTTTTC  
 CACAGCTCGG AACTGAGAG GTACAGCTCG TCGAGCGCGG GTCTACTGCG TCTGCTGCGG TCTGCTGCGG TCTGCTGCGG TCTGCTGCGG TCTGCTGCGG  
 4401 TGTGCGCGCG CGAGACCGAA GAGCTGTGGA CTGTTGGGAC ACGGCTGCGG GCGGCTGCTG ACACCGCTTC TCGATATCTT TGTCTGTGTA CTCTAACCCA  
 ACACCGCGCG GCTCTGCGCT CTGAGACCTT GAGAACCTCT TCGCGCTGCG GCGGCTGCTG GCGGCTGCTT TGTGCGGAGG AGTGTAGAA ACAGGACAAAT GAGTTGTGCT  
 4501 ACTTCTGCGG ATGATGCTGT GACTGCGTGG GTAGGCTGCG GTGCGGCTGCG GTGCGGCTGCG CACTTTTGGC GTCTTACTGT AGCTGCGGAG GCGAGTGTGCG  
 TGAGAGCGCG CTGAGCGCGC CAGTCCACCG CAGTCCACCG CAGTCCACCG CAGTCCACCG CAGTCCACCG CAGTCCACCG CAGTCCACCG CAGTCCACCG  
 4601 GTGTGAGAGG GGTGTGAGGT CGAGCTGCGA TCGACCTTCA GAGAGACCGA ACGAGGAAAT GAGAGCAGCG TCTGTGCTCT CTCTTCCCGG AGCCACCGAT  
 CCGAGTCCCA CCGAGCTTCA GCTGAGCGGT AGTGTGAGGT CTCTCTGCGT TCTGCTCTTA CTGTGCTGCG AGTGTAGAA ACAGGACAAAT GAGTTGTGCT  
 4701 CACCGCTCAA AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT  
 GGTGAGAGGT CCGAGCTTCC ACTGTGCTTA TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG  
 4801 GAGGAAATTA GCGCGGAGCT GAGGAGAGTC GATGCGCGCT TTGGGAGTCT GCTAGCGCGA CTCTGCGCTT AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT  
 CTCTGCTTCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT GCGCGGAGCT  
 4901 TCAATCTTTC CCGCTGCGGA AACTGCGCTT AGGTGTGATTT TTTTCTGCTG TCTGCTGAGG AGCGGCTTCT CACTCTAGCG GCGGCTGCGC CCGTGTGCGG  
 AGTCAAGAGG GTGAGCGCTT TTGAGCGAGG TCGAGAGAGG AAGAGAGGAG AGAGAGAGGAG AGAGAGAGGAG AGAGAGAGGAG AGAGAGAGGAG AGAGAGAGGAG  
 5001 TATATCTGTA AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT  
 ATATAGAGAT ACAGGAGAGG TCGAGAGAGG GTGTGCGGCT GCGGAGAGCG CCGGCTTCTT GCGGCTTCTT GCGGCTTCTT GCGGCTTCTT GCGGCTTCTT  
 5101 CCACTCTGCT GAGAGCGCTT ATACTGTGAA TATGAGCTTT GATGCTGCTG GCGGCGGAGT GCGGCTTCTT GCGGCTTCTT GCGGCTTCTT GCGGCTTCTT  
 GTTCAAGAGG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 5201 CCGTCTGCTC ACTTCTGCTT TCGGCTGCGT TCGAGAGAGG TATGAGCTTT TCGAGAGAGG TCGAGAGAGG TCGAGAGAGG TCGAGAGAGG TCGAGAGAGG  
 GTTCAAGAGG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 5301 TTTTCTGCTT AGTCTCTGCT CCGTCTGCTT CCGTCTGCTT CCGTCTGCTT CCGTCTGCTT CCGTCTGCTT CCGTCTGCTT CCGTCTGCTT CCGTCTGCTT  
 AAGAGAGGAG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 5401 GAGGCTTCTA GCGGCTGAGC TTTGCGCTCC CTATCTGCTA TATTTGAGT CCGGCTTCTA TTTTCTGCTT TTTTCTGCTT TTTTCTGCTT TTTTCTGCTT  
 GTTCAAGAGG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 5501 ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC ATCGCTGCTC  
 TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG TCGGCTGAGG  
 5601 ATATGCTGCT AGATGCGCGC TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT TCGAGAGAGG AGCTCTGCTT  
 TATGAGAGG TCGAGAGAGG ATGCGGCGCA GCGGCTGCTT GCGGCTGCTT GCGGCTGCTT GCGGCTGCTT GCGGCTGCTT GCGGCTGCTT GCGGCTGCTT  
 5701 GCGGCTGCTA TTTATCTCTT TCGGCTGCTT CAAAGAGGAG TATGAGCTT TATGAGCTT TATGAGCTT TATGAGCTT TATGAGCTT TATGAGCTT  
 GCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT TCGGCTGCTT  
 5801 GTGCTGCTT GGTGCTGCTT GAGCTGCTT GTTCTGAGT GTGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT  
 CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT  
 5901 TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT TGTGCTGCTT  
 AAGAGAGGAG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 6001 CAGGCTGCTT CAGGCTGCTT TATGCTGCTT CTGCGCTGCT TCTGCTGCTT CTGCTGCTT CTGCTGCTT CTGCTGCTT CTGCTGCTT CTGCTGCTT  
 GTTCAAGAGG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 6101 GTGCTGCTT TTTTCTGCTT CTATGCTGCT GAGGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT GTGCTGCTT  
 CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT  
 6201 TAGGCTGCTT CAGGCTGCTT TTTTCTGCTT TTTTCTGCTT GTTCTGCTT GTTCTGCTT GTTCTGCTT GTTCTGCTT GTTCTGCTT GTTCTGCTT  
 ATCGGCTGCTT GTTCTGCTT AAGAGAGGAG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 6301 CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT CCGCTGCTT  
 GAGAGAGGAG GTTCAAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG ATCTGAGAGG  
 6401 GCGCTGCTT CAGGAGAGG GAGTGTGCTT TCGTAGGAG GTCTGCTGCT CACTGCTGCT GAGGCTGCTT GAGTCAAGC CTACAGAGG ACTGAGAGG  
 CCGAGAGGAG GTCTGCTT CCGAGAGCTA AGATGCTTCT TCGTAGGAG GAGAGAGGAG CAGGCTGCTT GAGTCAAGC CTACAGAGG ACTGAGAGG  
 6501 AGTGTGCTT TCGTAGGAG TCGGCTGCTT CCGAGAGCTT CCGAGAGCTT GAGGAGAGG GAGGAGAGG GAGGAGAGG GAGGAGAGG GAGGAGAGG  
 TCGCTGCTT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT AGATGCTTCT  
 6601 TCGGAGAGT TCGTAGGAG AAGAGAGT ATCTGCTT GGTGTGCTT GAGTCTGCT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT  
 ACCCTGCTT AGATGCTTCT TTTTCTTAA TATGAGAGG GTTGTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT  
 6701 GAGGAGAGT CAGGCTGCTT GGTGTGCTT AGGAGAGT CCGGCTGCTT TCGTAGGAG TCGTAGGAG TCGTAGGAG TCGTAGGAG TCGTAGGAG TCGTAGGAG  
 CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT  
 6801 GAGGCTGCTT TCGAGAGG GAGGCTGCTT TCGTAGGAG GTCTGCTGCT CACTGCTGCT GAGGCTGCTT GAGTCAAGC CTACAGAGG ACTGAGAGG  
 CCGGAGAGG GTCTGCTT CCGAGAGCTT TCGTAGGAG GAGGAGAGG GAGGAGAGG GAGGAGAGG GAGGAGAGG GAGGAGAGG GAGGAGAGG GAGGAGAGG  
 6901 ATCGAGAGT CAGGCTGCTT GTTGTGAGG ACTGAGAGG CCGAGAGG TTTGAGGAG GTTGTGAGG GTTGTGAGG GTTGTGAGG GTTGTGAGG GTTGTGAGG  
 TCGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT CAGGCTGCTT  
 7001 CTGAGAGG CCGTGTGCTT TCGAGAGG ATCGAGAGT GAGGAGAGG GAGGAGAGG ATCGAGAGG CAGTGTGAGG CAGGAGAGG TTTCTGCTT  
 GAGTGTGCTT CCGAGCTTCTA AGTGTGCTT TCGAGAGG GAGTGTGCTT TCGAGAGG GAGTGTGCTT TCGAGAGG GAGTGTGCTT TCGAGAGG GAGTGTGCTT  
 7101 TATGAGAGT CCGTGTGAGT GATGAGTCT TCGAGAGG GAGTGTGCTT TCGAGAGG GAGTGTGCTT TCGAGAGG GAGTGTGCTT TCGAGAGG GAGTGTGCTT

FIG. 4B-2



[illegible]

**FIG. 4B-3**

```

10801 GTCTGGAAA AAAAAGAGA GAGAGGAGG TGAAGCCCA ATAACTTAA CATTTCTGT GTTGTCTTG CTTAGTCEA TTCTGTAAQ CAATCTGGC
CAGAGCTTTT TTTTCTCT CTCTCCCTC ACTCTGGGT TATTGAATT GTAAAGACAC GACATCAGT AAGACATTC GTTACGACG
10901 TTGCTCCAA GTTGGAGAT AACTTCTT TTTAAAGGT ATTGACTG CTTATTTT CTGTTTAT TATGCTCTG AGATGGAC CAGGACCT
AAGGAGGTT CCATCTCA TGTAAAGA ATATTTTCA TAAAGAGAC GAATATAAA GACAAATAA ATACACGAC TCTACCTTG GTTCTGGA
11001 TGGCAGCAA GCTTACTGT TTAACACTA GCAATCTCC AACTTGCAC TGGGGATC TGGCAGGG TTCTACACT GAGCCACT CCCCACCC
AAGGTTGTT CCAATGACA AATGCTACT CGGTATGAG TCGAAGCTG ACCGCTAG ATCGGTTCC AGATGTTGA CTGCTGTGA GGGGTGGGG
11101 ATCCCTCTT GAGATCTT AAGCACTC ATACCTAGC TTGATCTT TAAAGCTC TTAATAGG TCGTT
TAGGAGAGA CTTCTAGA TCGTCAGG TATGATGG AACTAGAA ATCTGCGG ATGATCTG ATCAA

```

FIG. 4B-4




FIG. 5A






FIG. 5B

<b>VEGF project</b>	<b>VEGFR2 project</b>	<b>Tie2 project</b>
<b>Screening primers</b>	<b>Screening primers</b>	<b>Screening primers</b>
Primers: VF1-VR1A Product size: 0.69Kb	Primers: KF1-KR1 Product size: 0.45Kb	Primers: TF3-TR1 Product size: 0.45Kb
<b>PCR program</b>	<b>PCR program</b>	<b>PCR program</b>
Hot start	Hot start	Hot start
94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec
40 cycles	40 cycles	40 cycles
<b>Confirmation primers</b>	<b>Confirmation primers</b>	<b>Confirmation primers</b>
Primers: VF2-VR2 Product size: 0.98Kb	Primers: KF2-KR2 Product size: 0.58Kb	Primers: TF2-TR1 Product size: 0.47Kb
<b>PCR program</b>	<b>PCR program</b>	<b>PCR program</b>
Hot start	Hot start	Hot start
94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec
40 cycles	40 cycles	40 cycles

FIG. 6

**FIG.7**

*FIG.8*





FIG.9



**pTKLG-Fos/VEGFR2  
targeted transgenic vector  
(Yellow-green luciferase)**

**pTKLR-Vn/VEGF  
targeted transgenic vector  
(Red luciferase)**



**FIG.10**

## SEQUENCE LISTING

<110> XENOGEN CORPORATION

<120> TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED  
THEREWITH

<130> PXE-008.PC

<140>

<141> 1999-12-16

<150> 60/152,522

<151> 1999-09-03

<160> 39

<170> PatentIn Ver. 2.0

<210> 1

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer PGKF

<400> 1

atcgaattct accgggtagg ggagggcgtt t

31

<210> 2

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer PGKR

<400> 2

ggctgcaggt cgaaaggccc ggagatgagg

30

<210> 3

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NeoF

<400> 3

acctgcagcc aatatgggat cggccattga ac

32

<210> 4

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NeoR

<400> 4

ggatccgcgg ccgccccag ctggttcttt ccgcctc

37

<210> 5  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer TKF

<400> 5  
ggatcctcta gagtcgagca gtgtgggtttt 30

<210> 6  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer TKR

<400> 6  
gagctcccgt agtcaggttt agttcggtccg 30

<210> 7  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F5R51

<400> 7  
gtacatttaa atcctgcagg 20

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F5R52

<400> 8  
agctcctgca ggatttaa 20

<210> 9  
<211> 77  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F3R31

<400> 9  
ggcccgggct taattaatgc atcatatggt accgtttaaa cgcgccgca agcttgctga 60  
cgcgcgccg gccggcc 77

<210> 10  
<211> 77  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer F3R32

&lt;400&gt; 10

gatcgggccgg ccggcgcgcc gtcgacaagc ttgcggccgc gtttaaaccg taccatatga 60  
tgcattaatt aagcccg 77

&lt;210&gt; 11

&lt;211&gt; 39

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN1R

&lt;400&gt; 11

ctgtatttaa atctgcccac cctattcagg acagtagtc 39

&lt;210&gt; 12

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN1F

&lt;400&gt; 12

ccaatgcatc aaccagcca ggaggagtgc g 31

&lt;210&gt; 13

&lt;211&gt; 38

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN2R

&lt;400&gt; 13

aacgcgtcga cttcgagat gtttcgggga taaccagg 38

&lt;210&gt; 14

&lt;211&gt; 40

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN2F

&lt;400&gt; 14

ttggcgcgcc ccatagagaa gagacaccaa aggcacgctc 40

&lt;210&gt; 15

&lt;211&gt; 36

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer FosB1F

&lt;400&gt; 15

ctgtatttaa atcccgtttc tcactgtgcc tgtgtc 36

<210> 16  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer FosB1R

<400> 16  
gtctcctgca ggcttcctcc tcttggtcc ttgcg 35

<210> 17  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer FosB2F

<400> 17  
aacgcgtcga cggatgggat tgacccccag ccctc 35

<210> 18  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer FosB2R

<400> 18  
ttggcgcgcc ccttgccctcc acctctcaaa tgc 33

<210> 19  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VF1

<400> 19  
acctcactct cctgtctccc ctgattccca a 31

<210> 20  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VR1A

<400> 20  
gctctggcgg tcacccccaa aagca 25

<210> 21  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VF2

<400> 21  
ccctttccaa gaccggtgcc atttgagc 28

<210> 22  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VR2

<400> 22  
actttgcccc tgtccctctc tctgttcgc 29

<210> 23  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KF1

<400> 23  
gctgcgtcca gatttgctct cagatgcg 28

<210> 24  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KR1

<400> 24  
ttctcaggca cagactcctt ctccgtccct 30

<210> 25  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KF2

<400> 25  
cagatggacg agaaaacagt agaggcgttg gc 32

<210> 26  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KR2

<400> 26  
gaggactcag ggcagaaaga gagcg 25

<210> 27  
<211> 29  
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TF3

<400> 27

agcttagcct gcaaggggtgg tcctcatcg

29

<210> 28

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TF2

<400> 28

caaattgcacc ccagagaaca gcttagcctg c

31

<210> 29

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TR1

<400> 29

gctttcaaca actcacaact ttgcgacttc ccg

33

<210> 30

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F51

<400> 30

cccagtgctc ctgatttagg gagagcacct gag

33

<210> 31

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer R51

<400> 31

ccagactgcc ttgggaaaag cgcctc

26

<210> 32

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F52

<400> 32

cagtgaagt cttctctgtc cctcaatcgg ttctg

35

<210> 33  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer R52

<400> 33  
tggatgtgga atgtgtgcga ggccag

26

<210> 34  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F31

<400> 34  
aatcaaagag gcgaactgtg tgtgagaggt cc

32

<210> 35  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer R31

<400> 35  
cggctcccca aaatgtggaa gcaagc

26

<210> 36  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F32

<400> 36  
gaatccatct tgctccaaca cccaacatc

30

<210> 37  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer R32

<400> 37  
cgctcctct cccagtcctc cccttg

26

<210> 38  
<211> 4998  
<212> DNA  
<213> Mus sp.

<400> 38



tccaccacc	tgtttctcac	gtccccggcc	ttcctagtta	acttcatggt	taaagaagcc	60
tcacccggg	aggggtgtgt	gccacagaag	gaaggggtgt	cccacaagcc	cccagtgtct	120
ctgatttagg	gagagcacct	gagcccagtg	agagtcttct	ctgtccctca	atcggttctg	180
aaattcccca	cttgccctcc	ttatccaggg	gacagggctg	cccaccctat	tcaggacagt	240
agtcttaaac	tcgtagccaa	cagacttttt	attgggctgg	gagaaagaga	tgaggctcct	300
gaagctcagc	cgagtgggct	ctgattccta	cttctcagag	gtcgggcagc	ccagccaata	360
ctgagcaatg	gagcgtgggt	agggaggatt	cacagagtcc	actcgccggg	ttctaagggt	420
gactcggtag	tatttgtctg	aaagaaagaa	tggaaaaagg	gttatgtgag	attctgcctg	480
atcctgtcca	ctgggtccca	gaaggataaa	ggctttttct	cagaagggaa	agtgaacatc	540
caccaagcag	ataatgtcac	catctacagg	ctgtgttcag	caccagggga	ccaagacctg	600
caggcaaggc	ctagccaaaa	ccagtctaag	gagtagaaag	gggctccac	ctccagagaa	660
gaaatagacg	ctctgaatgg	gctcgcaggt	ggcaggtaga	agccagtcca	tatcataatc	720
atagttgttg	taggttccta	gcccactctc	ctcgtcggag	aacaaagaga	accagattga	780
acgtgatgaa	cgacgggagt	tcgagctctg	gctgcgtctg	tggccacgcc	ctcggcgtga	840
acgatagcgc	tttcggcttc	tacgcttaga	cttctgtttt	ttggcttggg	cagagtggga	900
taaggagcca	gtgacgtaga	tgcggccggc	catagcagcg	tccactttcc	ctggcacacc	960
atgccagttc	cggctgatga	attggggttc	tctggctcca	tctgtaacag	ggaaggggtt	1020
aatgcacttg	cgagattctg	gctttgattt	ctccagcaag	gttgtctgtc	tatctattta	1080
tctatcttta	tctatgtatc	tatctatata	tctatgtatc	tatctatcta	tcatctacct	1140
acctacttac	ctatctatgt	atctatctat	ctatcatcta	cctacctact	tacctatcta	1200
cctattttatt	tgtttggttg	ttttctttga	aacaggatct	tagcacctac	ctatggctgg	1260
tttgcaactc	actatgaagc	cataactggc	ctcttaactc	acaaagatcc	acttgctgtg	1320
gtctctgagt	gctgggatta	aaagcatgtg	ccactacacc	cagctccagt	aggaccttta	1380
gaacacattt	gctatgcctt	gcctaagaca	cacaactcag	tccccaggcc	ccagcctccc	1440
tgtctagagc	tttttcccat	cctctctcca	ctgtatccct	tgaatctctg	ccccatccga	1500
aaccctcag	cgcgagccc	ctccttctgc	tgtgttaggc	aaagtccaag	gtatgggata	1560
caaataagagc	caagcctcat	cccccaaaag	tcaacagaag	caaagtctag	ccagagcaaa	1620
cagctcttga	tcagtgggtg	cacagttcca	ggccccctcc	ctggaagccc	ccactatcac	1680
agcccagttt	ccagagaaag	aagccagcct	tgtctccct	ccataccaga	ggatctgccc	1740
cagaagagga	gttcgaaaaat	gttctcccag	ctgtcccgt	gaagcaaggc	aaagtgtctc	1800
aacacggctg	acagagagct	gccttcgcac	tctcctggc	tgggttgctg	ctgaaattcg	1860
tactcccagt	actgcttccc	tgaggagcag	aacagctggc	atcaggagag	atctgaccaa	1920
ggcagagagg	aatcatggaa	tagaacaggg	actccaccac	ctgccccctt	ctcctccacc	1980
ctgagtaccc	ttgaagaagt	agacccttcc	ccggccactg	taacgggtgg	caggaagggc	2040
gaacgctgca	tcaacattgt	ctggtatgcc	actgaagcct	tcggagatgt	ttcggggata	2100
accaggggcc	aggaccccat	cctcaaagcg	ccagtactga	ctaccctgaa	agacagagat	2160
cagaaggggtg	aggacatacc	gctggccaca	gaagcagtc	tatatcctaa	actggctgtc	2220
acctgctcct	ggagctccct	actgctttgt	cttcacagct	ccccagcacg	tccatggcac	2280
cctttacctt	gcctcagact	taggtctggt	accttgaaca	agtaggtctt	cccctgacag	2340
ttgatgcgag	tgaaggcagc	atcgatgggg	ccctcaatgc	cccagacatc	ttggataagt	2400
ttgggggtacc	caggcctcac	tgccgtctca	tctagctcat	agcagtactg	ccctagaaca	2460
ggggaaactg	tgtgagaagc	agatgagcct	aaggcagatc	cgaccgccac	cagacctgtc	2520
catagagtca	cctcggaagg	caaagagggg	ccattctctg	agatccgtga	aggcgtcaaa	2580
gggctttcca	ctgcacagtt	cttccctctg	aaactcaggg	gtcccttgat	cagtgggtgtc	2640
gggccttagg	atctcctcct	gttgctccac	tttaggcgt	gggggtgctg	gctgttcctc	2700
aggatctagg	aaggctgtcg	gcttttagagt	gccgtccgtc	cgaggattta	ggtcaccggg	2760
tggagagggtg	ttctcgggtt	gcacaccggg	gttggtattg	ttcttgggct	cctccacgta	2820
gtcatagctc	caataatcat	cctctggcat	agtgaacacg	tccccccg	ttactgcagg	2880
cagaacgggg	agcagtgagt	gtcaggctgt	ggagggagcc	ccaggcccac	ccaccagggc	2940
tctgaactca	ccttgggggt	tgcactgtct	catgtagctg	gcacagcagc	tctgatagta	3000
agtgcaaaagc	tcgtcacact	gacacttctt	gctggccatg	aaacctgag	tgcagcggcc	3060
cctgcatgac	tctatgggag	ggaatatcag	gtttacagcc	caatctaggg	cacctgcccc	3120
acctgcactt	ccctaggtac	ccaccaatcc	cctcccacac	cttggtcagc	cagagaaacc	3180
catgccacca	gggctagtat	gaaaaagggc	ctcaggggtg	ccatggcagg	cctctagccc	3240
agggccttgg	caagctgggc	gcggagcttc	tggaaatctg	ctgtcctgcc	tgaaaaaaga	3300
agcagactga	agaagagttc	ctagttccct	gggtttctgc	cctttatttg	ctcatcctct	3360
ggcccagccc	cattgccttc	ctccaaacac	agctgcagca	aagggtcaca	ttcccagaac	3420
cccagcccca	ggagagctgg	gaaacagaaa	accctcgcca	agaccaaagt	cagttagggtc	3480
acgggcagga	gggataaac	gcttagctta	ctgggggagg	tggaaagaag	catgtgttgt	3540
cacctcttga	gccagtccc	ttaatctccc	tgagccttac	tttttataaa	gtgggaccat	3600
gggtgccttgc	ctcatcaggt	gttgagagat	tccgtgagct	agaacagaca	aaacgtttcg	3660
tgcctggagt	agcttccaac	tcatcccat	aagccgttat	cgatttactg	tttgatcagg	3720

ctaggtgctt gtcccatcct acccccgcgt tcgaatctgg atttttgggg caagaagggg 3780  
 ggttggggga gagctggcaa gcactttggg ggaggttttc ttttcttctc ataaaagaac 3840  
 aaagcttcat ttctggcctc tccttggtct ctctaagctg ggtgttacag cataggaagt 3900  
 agtgggtcag agtctattct tctttcttta ttttttttag atttatttat tttatgtttt 3960  
 gtgtataagt gtctgtcac atgtgcatt gtgcaccaca tgcattgtct gtgtctatgg 4020  
 aggtcagaag agggctttga ataccctgga actggagttt tgaacagtta tgagctgccg 4080  
 tgtggatgct gagaatcaaa cccaggtcct ctgtaagaac aagtactctt aaaggctgag 4140  
 ccatctttcc agtcccagag cccattcctg aggcctttcac taatccattg atcctcgggg 4200  
 gaccaccctg gccacacctt caatgacctc atttatttta aaaaaaaat' ggactcattg 4260  
 ggcatacttt ctagactcac atactaagtg ggatttctct ataaagaagt gctcactggg 4320  
 gtagagtgcc aggttttggg ccaaattcca agcactggca cacttctgaa gccctccgt 4380  
 tttctgttct gtaatcacag gcgagcgtgc ctttgggtgc tcttctctat ggaccgag 4440  
 agtctcagcg gcaaatgaa acactaaatt ttactcccta cagacgcgtg aagcctaagt 4500  
 ggaaaccggc attaaagggc tttaagaatc tcaactgcga ttctttaacc atccggaggg 4560  
 gacgtggata catgtagcca gcttgcttcc acattttggg gagccgagcg agcggtagga 4620  
 aatggaagac agctctttac agccctttct acagcatctt gcacaccacc aaggggagac 4680  
 tggggagagg agggcgagcc aggtgtgggc gtggctggag acctggggta ggcttgcgcc 4740  
 tgcgtcgggg gcggagcccg tgaaacctag aggcggggcg tcaaatcctt gactctgctg 4800  
 ctacagggcg tgggtgctgt tgagcatctt agctccgctg tgcttagatt ggagcagcg 4860  
 tttgttccgg gcaccggcgt ctctaccctc ccgcgtctgg tccatgcttc tctctccctt 4920  
 catgcccttc ctaagtcgct gagtcccgga gctgcctcc tccttctgct tctacacttg 4980  
 tagccagca cctttacc 4998

&lt;210&gt; 39

&lt;211&gt; 11176

&lt;212&gt; DNA

&lt;213&gt; Mus sp.

&lt;400&gt; 39

gcagctgggc aaacgttggc gatgccggtg caaagtatat acccggtggt tagcagaagc 60  
 tgagaacttt tagccgaaag ccggtccctc aagccgaagc taggcaagta ggggaagaaa 120  
 aagaaaaaaa aaattccaga gaagcttcca gagcctcctc ctcttccctc ttccttcaaa 180  
 aggactgcaa gtccgcagtc accctccacc cagcaagagt tagggcctcg aaccccggtc 240  
 acgctgcctc cgctcctgc cgaacgtaac gggggaccog tgcgtaaaagc gtgacgcgtc 300  
 ggaatcctcc gtctgacgog gggcacgcac aggcgcgagc cctccgccc gccccgcccc 360  
 tgacgtccgg gcacgttcta ttttggaacg ccgagggccac gttgctaagg gagggggcag 420  
 cgtggctttg tgatttctg tcgcgcgagc ctttagccaa tcagcgttcc ctctctattt 480  
 gtagagcgtg gctcccttcc ttgctttttg tggttcttcc cgtgctgggg gtctccaaga 540  
 ggagagctag gattcttgtc gcgatcggga ctgctgtgca ccccatgggtc tgcgaggact 600  
 tgtgtggacc tgggtctgtg tcataagcta gaggcttttg gctgagtgtt agcgctcta 660  
 agggggaact gaaggcctca tcttctcag gcacacatat acgtgctcct gagctctaga 720  
 cactcagtc ttccgaggtg ttcaaact agatgagcta gcctacggag aggcagccag 780  
 gtggtctcta aaaggtctgc ctcccttag ttcccaggct ctgattggcc agggattcag 840  
 ccttccctc gccacgcccc ctagagtagt taagcctcta ggattccact tgcgggaagg 900  
 gggggggggg gggcgtgatg gacgttctt ggggacgcag atcctatgtc accccatccc 960  
 ctgcaagaca gtctgagaga ttctcgctgt cacttttctc tgccatcag ttactgaaa 1020  
 cctgtcagtc tactgggaa gagacagaca ctcggaaggg atgctctcaa ctctaggcc 1080  
 ggtcccccaa caccgttgga actgggatct ccgctgcgg gagccctcat gcagtggggg 1140  
 gtgtgtttgt gtgtgagtg agaggaaggc ttggctaagg cctctccctc tccctccctc 1200  
 tgtgtggggg gttgggggt tttggctgta tgtgtgtgtg aatgtctgtg gctccatccc 1260  
 gggagtttgt caccaggttc tgtccagcct cctctccac ccaccccccc acacctaga 1320  
 gtcaccaacc cgggtgtgta ttcaccacc gctggaaccg tgcaaccttt ccccgaggaa 1380  
 gaaggaggag gtagaaggca gttgaacaga atctctcatt aacctgog tcacgggtga 1440  
 gtggaagggt ggggtgtgtg gctttttgcc tgtgacacac acatccacac ccgctcacc 1500  
 tgtgtcact cacagggtcg gtctctotta tctctctgg gcgtgtgtgt gtcgggtggt 1560  
 ttgtttgtgt gtctacgct gtgtgtgtat gtctcaccg taggagtgcc gccggtctcg 1620  
 gggaaatgcc cggctccttc gtgccaacgg tcaccgcaat cacaaccagc caggatcttc 1680  
 agtggctcgt gcaaccacc ctcatctctt ccatggccca gtcccagggg cagccactgg 1740  
 cctccagcc tccagctgtt gaccttatg acatgccagg aaccagctac tcaacccag 1800  
 gcctgagtg ctacagcact ggcggggcaa gcggaagtgg tgggccttca accagcaca 1860  
 ccaccagtg acctgtgtct gcccgccag ccagagccag gcctagaaga ccccgagaag 1920  
 agacagtaag tatgagcct caggagttgg gatggaggag cctagctagg gatgtgggt 1980

cagtttgtac	agtgccttgc	tgccatgcat	gaagatccct	agcacagcat	aagccaggag	2040
tggttatgca	gacctgtaac	cccagctctc	agaaggtgga	ggcaggagga	gcaggagttc	2100
gaggccagcc	tgtgctactt	atggagtcca	gcctgcactg	caagagatca	ttattttcaa	2160
aagttggcct	tggggggagg	tgggtgaggg	aagtaagaga	aagtgcaggt	aatattgtca	2220
cttaatagtt	ggaggttctt	ctgaggcctc	aagtctgaag	gaactttacc	attctggcca	2280
gtgaggagta	gggggttatta	tttgggggtt	aggaggaagg	aagttttctt	agggctgata	2340
gaggtacccc	cagatctcat	ggtccttatc	tctgactcag	cttaccctcag	aagaagaaga	2400
aaagcgaagg	gttcgcagag	agcggaaacaa	gctggctgca	gctaagtgcg	ggaaccgtcg	2460
gagggagctg	acagatcgac	ttcaggcgggt	aaggaggagt	ctgggggtgt	cttgaggccg	2520
tgtggggagc	actctgcctt	gttcttcccc	cgtttctcac	tgtgcctgtg	tcctaaacga	2580
ggaaaccccc	tcttagggaa	caggggtcag	tataggctga	tggagtggct	ccatatgcat	2640
gctcagaccc	atgcccactt	actttcgact	gttccccact	ttccctgaat	atgtccccac	2700
atgtcacctt	cctggctttc	tctcagccta	aggagacaag	ctagaggagg	taattctctc	2760
accttctttt	cttactctaa	taataatcca	ttttgccttc	ctgcctccat	ttttttttcc	2820
tgagctgggg	atctacctgt	cgtagttcag	ccctcctccc	ccaacttgat	agcctcaagt	2880
ttcagccctt	ggctgagatg	ccatcatcct	gactggctct	ggctggaaac	tattttgtgc	2940
taagtcaatt	ccttgtctgc	tacttcagct	atctacagty	ctgccgaact	tgagctgggtg	3000
gcgcccacca	agcccacttc	tttctctctt	ttttacctca	gtgcaacccc	ccacacacaa	3060
aacttcatgc	ctgccccttg	aaaccagggt	gcgtctctga	ctccccgtcg	ggaggctgaa	3120
ggagatgggt	aacagaacct	cattaaaaac	aacacataag	cattacctac	tgactcaaca	3180
aactgtagt	tttttctttt	ttcctctcaa	aaaattatct	cgtttgttta	tttattatct	3240
gcttatgttt	gagtgagtgc	tgggtgcacca	cagcacacat	acgaggctcag	agggaaatct	3300
tcatagtttg	ttctctcctt	ccgtgttgtg	ggtgcttgct	ggcaatctcc	ttcactcagt	3360
gagctacaat	gcccccttct	gccctttaag	gcagagtact	ccttagtaca	gggggacctc	3420
ttcctcgggc	tctcaaagtt	gagattacaa	atgttcacca	tcacaccagg	cttgaggttc	3480
ttgcctatca	gtgacgtcca	ctcctgccta	gcttcttccc	aaccatcttt	tagtctgatg	3540
gggaaaccga	ggcacagagta	gcatgggtct	ccaggatttc	ctcttagggg	acgggtccct	3600
cagttgggag	ggagctgtcc	agccccctg	atcagcagca	agaatgtatg	agtgtggggg	3660
tgggcgggtg	aagctactct	gtgtggctcg	tgaccagcaa	ttctcctttc	tctgtctcct	3720
atgacctggc	cctgtctggg	tccattagga	aactgatcag	cttgaagagg	aaaaggcaga	3780
gctggagtcg	gagatcgccg	agctgcaaaa	agagaaggaa	cgcttgagg	ttgtcctggg	3840
ggcccacaaa	ccgggctgca	agatccccct	cgaagagggg	ccggggccag	gcccgcctgg	3900
cgaggtgaga	gatttgccag	ggtcaacatc	cgctaaggaa	gacggcttcg	gctgggtgct	3960
gcccggccct	ccaccaccgc	ccctgcccct	ccagagcagc	cgagacgcac	cccccaacct	4020
gacggcttct	ctctttacac	acagtgaagt	tcaagtcctc	ggcgaccctt	tcccctgtgt	4080
tagcccttcg	tacacttcc	cgtttgtcct	cacctgccc	gaggtctccg	cgttcgccgg	4140
cgcccaacgc	accagcggca	gcgagcagcc	gtccgaccgc	ctgaactcgc	cctcccttct	4200
tgctctgtaa	actctttaga	caaacaaaac	aaacaaaccc	gcaaggaaaca	aggaggagga	4260
agatgaggag	gagaggggag	gaagcagtc	gggggtgtgt	gtgtggaccc	tttgactctt	4320
ctgtctgacc	acctgcccgc	tctgccatcg	gacctgacgg	aaggacctcc	tttgtgtttt	4380
gtgctctgtc	tctggttttc	tgtgcccccg	cgagaccgga	gagctgggtga	ctttggggac	4440
aggggggtgg	gcggggatga	acacccctcc	tgcatactct	tgtcctgtta	cttcaaccca	4500
acttctgggg	atagatggct	gactgggtgg	gtaggggtgg	gtgcaacgcc	cacctttggc	4560
gtcttacgtg	aggctggagg	ggaaagagt	ctgagtgtgg	ggtgcagggt	gggttgagg	4620
cgagctggca	tgcacctcca	gagagaccca	acgaggaaat	gacagcaccc	tcctgtcctt	4680
cttttccccc	accaccccat	ccaccctcaa	gggtgcagg	tgaccaagat	agctctgttt	4740
tgctccctcg	ggccttagct	gattaactta	acatttccaa	gaggttacaa	cctcctcctg	4800
gacgaattga	gcccccgact	gagggaagtc	gatgccccct	ttgggagtct	gctaacccca	4860
cttcccgcgt	attccaaaat	gtgaacccct	atctgactgc	tcagtctttc	cctcctggga	4920
aaactggctc	aggttggtg	tttttccctg	tctgtacag	agccccctcc	caactcaggc	4980
cogctcccac	ccctgtgcag	tattatgcta	tgtccctctc	acctcacc	ccacccagg	5040
cgcccttggc	cgctcctggt	gggccttact	ggttttgggc	agcagggggc	gctgcgacgc	5100
ccatcttgct	ggagcgcttt	atactgtgaa	tgagtggctg	gattgctggg	cgcgccggat	5160
gggattgacc	cccagccctc	caaaactttt	cctgggcctc	cccttcttcc	acttgcttcc	5220
tccctccctt	tgacagggag	ttagactcga	aaggatgacc	acgacgcac	ccgggtggcct	5280
tcttgctcag	gccccagact	ttttctcttt	aagtccttcg	ccttccccag	cctaggacgc	5340
caacttctcc	ccaccctggg	agccccgc	cctctcacag	aggtcgaggc	aattttcaga	5400
gaagttttca	gggctgaggc	tttggctccc	ctatcctcga	tatttgaatc	cccaaatagt	5460
ttttggacta	gcatacttaa	gagggggctg	agttccact	atcccactcc	atccaattcc	5520
ttcagtccca	aagacgagtt	ctgtcccttc	cctccagctt	tcacctcggt	agaatccac	5580
gagtcagatt	tctattttct	aatattgggg	agatgggccc	taccgcccgt	cccccggtct	5640
gcatggaaca	ttccataccc	tgtcctgggg	cctagggttc	aaacctaata	ccaaacccca	5700

ccccagcta tttatccctt tccctgggtcc caaaaagcac ttatatctat tatgtataaa 5760  
taaataatatt atatattgagt gtgcgtgtgt gtgcgtgtgc gtgcgtgtgc gcgtgcgtgc 5820  
gagcttcctt gttttcaagt gtgctgtgga gttcaaaatc gcttctgagg atttgagtc 5880  
gactttctgg ctgtcccttt ttgtcacttt tttgttgttg tctcggctcc tctggctgtt 5940  
ggagacagtc ccggcctctc cctttatcct tctcaagtc tgtctcgctc agaccacttc 6000  
caacatgtct ccactctcaa tgactctgat ctccggtctg tctgttaatt ctggatttgt 6060  
cggggacatg caattttact tctgtaagta agtgtagctg ggtggtagat tttttacaat 6120  
ctatatcgtt gagaaattctg ggtggaaatg tctgatcagg agaagggcct gccactgccg 6180  
accacaattc attgactcca tagccctcac ccaggctgta tttgtgattt ttttcatttt 6240  
gtttttttgt attttgcacc tgaccccggt ggtgctgggg cagtctatca ctgggcagct 6300  
ccccccccc ccttggttct gcactgtcgc caataaaaaag cttttaaaaa actgtatcct 6360  
tcaggtcaaa gtgtctgttt tccctggaca tctactacat ggcttccttt cagaaaaacg 6420  
gagtttgat tgctaggga gtcttgctgg cacttagtgg gacgcctaac gaatcagaac 6480  
ctacaacggg actaaaaagga agtgagact tgctaggtt tcccatgttc ccaggctggg 6540  
ccacctactt gaaaaataa ggggcggaaa agtgtaaggt accaaatttg gtgaaggggtc 6600  
tgggagaatt tcatgatcgg aaaagaattt attcacttg ggtgtgcaat gaactttcag 6660  
caacagttaa gggcaagggt gtaaaagctg ggcacaactt gtaaatccta gcatttgaga 6720  
ggtgaggga cggggatcaa ctggtggagt tcagtgtcat gtggatcgta gataccaagc 6780  
gcaaagatct gctatgggga gagggcttgg tacaccaggg gagccagaag tttcgtgggtg 6840  
agggtagtgg agggcaagtg gagagtgaga gttagcctca gggagattct acaggcaatg 6900  
atgcagagtt cagacgctcc ctttgaaagc actagagagc cgcagcagggt tttgagcaga 6960  
gaaggtaga gttaggtggt ctcttctagc ccatcccagg ctgaggaggga cgtgagggt 7020  
ttcaagaagg atcgagaatg gaaagcagag gagaagaagg atccaagagg catggaggag 7080  
gcagaacaca tttctcttct ttaatagcaa gcctggaaag gataacttgc tgcaggagga 7140  
gatgctcacc agtcgggtgg tctaggggggt tcttggaana gagaaggcat ttgctcaagc 7200  
ctcgggtccc ccattctcgc tcttctgtca gcttgtcttc cattaagtgt gtgtctcaag 7260  
gccacctgct tcaggactcc ttgtgagagc acctctatg ctcgagttca ttaaaaaacac 7320  
aattgcctgg tccgctgctc tctccactgg ctgagttacc tcaaaagacc agggctaaag 7380  
gtgtgatcac aactctatcc ccattactgc tccaacgcag agacaggact gagccggagt 7440  
gaacaaatga acaaaaatga ctaataatgc atgcgtgatt aaatacataa aagagcagat 7500  
gactggatga gcaaatcgtt taaggagaga cagcaagatc ctagaatttt ggagactaat 7560  
ttaaatccat ctttgagatg catttggctg gaaattcctg ggaggaaaaa aagtgtaaat 7620  
atgaagagag aataaatgag aataggggtg gcttcagaga ggttaactgc gcgctggctg 7680  
cttttgtaga agaattgtga ttgcaggga caaaatggga tagatactcc cgcccgaaag 7740  
gtggaattga accactctgt cgctaaacag ctacagggtt gaagcctgca ccccgagca 7800  
ctgaggatca tccgggcgaa aggagctatt ttcagttagt tatataaagg cgagatacta 7860  
ctacttttta cacttatggt cattatttgt ggtatacagt agataattaa tttcaatggt 7920  
ttcgaacatt ttttttcaat ttttcttgtg aacatgtgtt tccctagtaa agtgttccgt 7980  
gaatgactct actaaataa aagtaagtag cttcatttgc atagcgcctt gcattttggg 8040  
aagcagcgcc taaagtgcct gtctccctaa ctaaaagcag aattttttgc aaagtgaana 8100  
gtcagtttta tttttgtttg tttgtttgct tgtttgttt taatggaaaa acttctcacg 8160  
cggcccatc gtagcagaat tgcagatttt ttgcaagcga gaagcaagac tttcgtaggg 8220  
tctgacggca cgcggccgca gagcgacacc tgccgttgc ttagagaact gcaagtatgt 8280  
agggaaatcta ctgagtcctt aggtgatgga gttgacaacc aactccctt gagtttagac 8340  
gctaaaaacc atcccttttt atatttatgt gattagccca gggaaactaa ggctcagaca 8400  
tggaataatc cacagccgag ttcttgtagc ccaactccct agtggcccta tggcaggag 8460  
ttgtgggttt aatatgctt gcccagggg gagaaagtgt tcaactggag gcgaggttt gaggtacgta 8580  
agcggagggt taccttgtta gatcctgggt gtctgcagaa cgtggtctcc ttctggctgc 8640  
tgctcaagtc aggtgtagaa ctctcagct cttctccagc accatgtctg cctgcttaat 8700  
gctttgcttc tttccatgac gataatgaac tgtgcctctg aaactgtaag tcagcccccc 8760  
agttacatgt tttcttttat aagagttgca tatatatatg tatgtatata tgtatgtata 8820  
tatgtatgta tatatatata tatatatata cagggtctca ctctttagct ctggctggcc 8880  
tgaaattcac tatgtagccc aggtatgctt gaactttgaa gcaatcttcc tgcctcagcc 8940  
tcccaatggt attacaggca tgagtcacaa caagccattt aaatcttatg atgacttata 9000  
agaagacaga aaatcagagt tcccttacct agttcacaga tccctacaat ctaacctcgt 9060  
tcgctccata aacagcccta cccaccctc ctggaactgc tttgaggaat gctgcagggt 9120  
ctcacaggca cactcctcct tgggttaatct cttcagcctg gttgccttcc ccccccattg 9180  
ccatgtggcc caaagcctct catcctgttc tcaaatacca ctgactagta aggtccccc 9240  
acctgacccg gtttaaatat tagaaaaggg tcaactttct cctgccacag acaaccaaac 9300  
caccatagtc ttgtcactta ctacctgact atgaagggt atagatgtct tcacaacctt 9360  
tctctgagcc tcagtttccc cacctgcata atgcatctga gacacagaat tccctagagc 9420

tgtggttctc	ctcattccta	gtgctgggac	cctttaatac	atttcctcat	gttgtggtga	9480
ccccaccacc	accataaaat	tatttccatt	gatacttcat	aactgtaatt	ttttctattg	9540
ttatgaatag	taatgtaagc	atttgtgttt	cccagtgatc	ttagatgacc	ctgtggaaga	9600
gtcattccac	cccaaagggg	tccccaccac	aagttaagaa	ttcctgccat	agaggaatca	9660
cagggaccat	ggattaacac	ttgggtcgac	ttttgggtcg	ccttctggga	ggcgctagag	9720
ctaatagcag	ctacatcaat	ttctgaaatt	ttgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	9780
gtgtgtgtgt	gccctgagtc	gggtgctgag	ataggccagt	ggcttttagt	ttcctggacc	9840
cattactcac	cagaactctc	ccctcacctg	attctttgat	gtgaacacta	tgtcttcata	9900
gtggcgggtg	caatagcagc	aacagtgaac	taaattttta	aagtagaact	cagctggaga	9960
tacaaatatt	gcagttttga	agttgggggt	gattgtctaa	taacttaata	acataaccca	10020
gaagagaggg	cccttggtct	tgcaaaactt	atatgcctca	gtacagggga	acgccagggc	10080
caagaagtgg	gagtgggtgg	gtaggggagc	aggggtgggg	gagggtatag	gggactttcc	10140
ggatagcatt	tgaaatgtaa	atgaagaaaa	tatctaataa	aaatttgaaa	aaaaatgtta	10200
ccccagtttg	gcctggatct	cactacctca	accagactgg	catgtgactc	tgctgagatc	10260
tgccacttct	tgccctcctg	gtgcagaaga	caatttttgg	aagttagttc	tcttcttcca	10320
tcttgtggat	tccagggatt	gaactcgggt	catcaggctt	ggctgcaagt	gacttactta	10380
gggtgtctccc	agaccctctc	ggtttgatta	gttagatgct	gcacttcatg	cctgactttc	10440
gcactatgta	gatagagcaa	tgtctataac	atctcctaca	atgatatgta	tatcaagagc	10500
caagtgatga	gatggctcag	tgggtaagag	cacagactgc	tcttccaaag	gtcccagagt	10560
caaatcccag	caatcacata	gtggcttcca	ttccctctta	tggaatgtct	gaagactgct	10620
acagtgtact	tacatataat	aaataaataa	atcttaaaaa	aaaaaaaccc	agccggggcgt	10680
gggtggcgac	gcctttaatc	ccagcacttg	ggaggcagag	gcaggcggtat	tcctgagttc	10740
gacgccagcc	tggtctacag	agtgagttcc	acgacagcca	gaactacaca	gagaaaccct	10800
gtctcgaaaa	aaaaaagaga	gagaggggaag	tgagagcgca	ataatcttaa	catttctgtg	10860
gttgtctttg	ctgtagtcta	ttctgataag	caatgctggc	ttgctcccaa	ggtaggaagt	10920
aacatttctt	tataaaaagg	atttgctctg	ctttattttt	ctgttttatt	tatgggtgctg	10980
aggatggaac	ccaggaccct	tggcaagcaa	ggctagctgt	ttaccaactga	gccatactcc	11040
agccttgcac	tgggggattc	taggcaaggg	ttctaccact	gagccacact	ccccaccccc	11100
atccctctct	ggaagattct	aggcagttcc	atacctagcc	tttgatcttt	taagacggtc	11160
ttactagagc	tcagtt					11176

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/90 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12N A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 30715 A (ISACOFF EHUD Y ;SIEGAL MICAH S (US); UNIV CALIFORNIA (US); CALIFOR) 16 July 1998 (1998-07-16) page 29, line 5 -page 30, line 10; claims	1-8, 10, 22-26, 32, 34, 35
X	WO 98 28971 A (LINK CHRISTOPHER ;UNIV TECHNOLOGY CORP (US)) 9 July 1998 (1998-07-09)	1, 2, 8, 10, 12, 14-17
Y	page 22, line 3 - line 6; figure 1  page 26, line 24 - line 28; claims 11-19; figure 1; example 3 page 18, line 9 - line 13  --- -/--	1, 2, 7, 10-12, 14, 17-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

26 June 2000

Date of mailing of the international search report

03/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Chambonnet, F

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CONTAG CH, ET AL.: "Visualizing gene expression in living mammals using a bioluminescent" PHOTOCHEMISTRY AND PHOTOBIOLOGY, (1997) VOL. 66, NO. 4, PP. 523-531. , XP000920570 the whole document	1,2,7, 10-12, 14,17-20
A	CONTAG PR, OLOMU IN, STEVENSON DK, CONTAG CH. : "Bioluminescent indicators in living mammals." NAT MED. , vol. 4, no. 2, February 1998 (1998-02), pages 245-247, XP000914588 the whole document	1
A	WO 98 36081 A (ANGELIS DINO A DE ;SLOAN KETTERING INST CANCER (US); MIESENBOCK GE) 20 August 1998 (1998-08-20) the whole document	1
A	WO 98 23633 A (CORNELL RES FOUNDATION INC) 4 June 1998 (1998-06-04) the whole document	1

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9830715	A	16-07-1998	AU	5090498 A	03-08-1998
WO 9828971	A	09-07-1998	AU EP	5619298 A 0948612 A	31-07-1998 13-10-1999
WO 9836081	A	20-08-1998	EP	0981633 A	01-03-2000
WO 9823633	A	04-06-1998	AU	7410398 A	22-06-1998